



Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd

181 - 205 Davy Road, Booragoon WA 6154,
Australia

Prepared by:

SLR Consulting Australia

Level 1, 500 Hay Street, Subiaco WA 6008,
Australia

SLR Project No.: 675.VX5890.00001

17 June 2024

Revision: 2.0

Revision Record

| Revision | Date | Prepared By | Checked By | Authorised By |
|----------|---------------|-----------------------|------------|---------------|
| 1.0 | 19 April 2024 | P. Dean S. Girando | E. Webb | E. Webb |
| 2.0 | 17 June 2024 | K. Griffith | E. Webb | E. Webb |

Basis of Report

This report has been prepared by SLR Consulting Australia (SLR) with all reasonable skill, care and diligence, and taking account of the timescale and resources allocated to it by agreement with Alcoa of Australia Ltd (the Client). Information reported herein is based on the interpretation of data collected, which has been accepted in good faith as being accurate and valid.

This report is for the exclusive use of the Client. No warranties or guarantees are expressed or should be inferred by any third parties. This report may not be relied upon by other parties without written consent from SLR.

SLR disclaims any responsibility to the Client and others in respect of any matters outside the agreed scope of the work.



Executive Summary

Alcoa of Australia Ltd commissioned SLR Consulting Australia Pty Ltd to undertake a fauna desktop assessment and targeted Chuditch (*Dasyurus geoffroii*), Quokka (*Setonix brachyurus*), and Woylie (*Bettongia penicillata*) survey for the Willowdale Mine Larego region as part of its Mining and Management Plan assessment by the Mining and Management Program Liaison Group. The survey was undertaken within a defined area that consists of approximately 35 individual polygons (the Survey Area) that cover 451.8 hectares and is located approximately 15 km southeast of the Waroona townsite, in the Jarrah Forest bioregion of Western Australia.

The objective of the survey was to identify key fauna values targeting Chuditch, Quokka, and Woylie within the Survey Area as part of the MMP assessment process.

The targeted fauna survey used a variety of detection methods including camera trapping, spotlighting, active searches, and opportunistic observations. Fauna habitat mapping was based on a combination of field observations, vegetation mapping, fauna habitat assessment data, and aerial imagery.

Two fauna habitats were mapped within the Survey Area, Jarrah-Marri Forest, and Drainage Line habitats. The Drainage Line habitat provides the most value for the targeted taxa as it is core habitat for the Chuditch, Quokka, and Woylie. The Jarrah-Marri Forest also provides value for targeted taxa as core habitat for the Chuditch and Woylie and supporting habitat for the Quokka.

One targeted taxon was recorded during the survey; secondary evidence of Chuditch was recorded in the form of scat within the Jarrah-Marri Forest. The Quokka and Woylie were not recorded during the current survey; however, they have previously been recorded within 200 m and 40 m of the Survey Area respectively.

In addition to the targeted fauna taxa, four significant taxa were recorded during the fauna survey:

- Forest Red-tailed Black Cockatoo (*Calyptorhynchus banksii naso*).
- Baudin's Cockatoo (*Zanda baudinii*).
- Western Brush Wallaby (*Notamacropus irma*).
- Rakali (*Hydromys chrysogaster*).

A further seven significant fauna taxa were assessed as having a high likelihood and six significant fauna taxa were assessed as having a medium likelihood, and 15 significant fauna taxa were assessed as having have a low likelihood of occurring within the Survey Area.

Three introduced taxa were recorded during the survey: Pig (*Sus scrofa*), Cat (*Felinus catus*) and Black Rat (*Rattus rattus*).



Table of Contents

| | |
|---|-----------|
| Basis of Report | i |
| Executive Summary | ii |
| Acronyms and Abbreviations | vi |
| 1.0 Introduction | 1 |
| 1.1 The Project..... | 1 |
| 1.2 Objective and Scope | 1 |
| 2.0 Background | 2 |
| 2.1 Statutory and Regulatory Framework | 2 |
| 2.2 Existing Environment..... | 2 |
| 2.2.1 Climate..... | 2 |
| 2.2.2 Interim Biogeographic Regionalisation of Australia..... | 3 |
| 2.2.3 Soil Landscape Mapping | 3 |
| 2.2.4 Hydrography..... | 4 |
| 2.2.5 Pre-European Vegetation | 5 |
| 2.2.6 Conservation Areas..... | 5 |
| 2.2.7 Environmentally Sensitive Areas | 5 |
| 2.2.8 Land Use..... | 5 |
| 3.0 Methods | 6 |
| 3.1 Desktop Assessment..... | 6 |
| 3.1.1 Literature Review | 6 |
| 3.1.2 Database Searches..... | 6 |
| 3.1.3 Likelihood of Occurrence..... | 7 |
| 3.2 Field Survey | 7 |
| 3.2.1 Survey Timing | 7 |
| 3.2.2 Field Personnel and Licences..... | 8 |
| 3.2.3 Habitat Assessment and Mapping | 8 |
| 3.2.4 Camera Traps | 8 |
| 3.2.5 Spotlighting | 8 |
| 3.2.6 Opportunistic Observations | 9 |
| 3.2.7 Active Searches | 9 |
| 3.3 Limitations..... | 9 |
| 4.0 Results | 11 |
| 4.1 Desktop Assessment..... | 11 |
| 4.2 Fauna Habitat..... | 11 |
| 4.3 Recorded Fauna..... | 13 |



| | | |
|------------|--|-----------|
| 4.4 | Significant Fauna..... | 16 |
| 5.0 | Discussion..... | 27 |
| 5.1 | Fauna Habitat..... | 27 |
| 5.2 | Significant Fauna..... | 27 |
| 5.2.1 | Recorded within the Survey Area | 27 |
| 5.2.2 | High likelihood of occurrence within the Survey Area | 29 |
| 5.2.3 | Medium likelihood of occurrence within the Survey Area | 31 |
| 6.0 | Conclusion..... | 33 |
| 7.0 | References..... | 34 |
| 8.0 | Feedback..... | 37 |

Tables in Text

| | | |
|-----------|--|----|
| Table 1: | Soil landscape subsystems and phases within the Survey Area | 4 |
| Table 2: | Hydrographic features in the vicinity of the Survey Area | 4 |
| Table 3: | Database search details | 6 |
| Table 4: | Likelihood of occurrence criteria | 7 |
| Table 5: | Survey timing..... | 7 |
| Table 6: | Field personnel..... | 8 |
| Table 7: | Limitations and constraints associated with the survey | 9 |
| Table 8: | Fauna habitats recorded within the Survey Area..... | 12 |
| Table 9: | Fauna recorded within the Survey Area | 14 |
| Table 10: | Significant fauna likelihood of occurrence within the Survey Area..... | 19 |

Graphs in Text

| | | |
|----------|---|---|
| Graph 1: | Climate summary of Dwellingup Weather Station | 3 |
|----------|---|---|

Figures in Text

| | | |
|-----------|--|----|
| Figure 1: | Chuditch scat recorded within Jarrah-Marri Forest (-32.9758445, 116.0860574)17 | |
| Figure 2: | Rakali recorded by camera trap 5890-CAM-07-EW within Drainage Line habitat (-32.9941287, 116.0524062)..... | 17 |
| Figure 3: | Western Brush Wallaby recorded by camera trap 5870-CAM-12-LB within Jarrah-Marri Forest (-32.97344506, 116.0826827)..... | 18 |



Maps

| | | |
|--------|--|-----|
| Map 1: | Survey Area..... | A-1 |
| Map 2: | Soil Landscape Mapping..... | A-2 |
| Map 3: | Hydrography..... | A-3 |
| Map 4: | Pre-European Vegetation | A-4 |
| Map 5: | Conservation Areas and ESAs..... | A-5 |
| Map 6: | Previous Surveys in the Locality | A-6 |
| Map 7: | Survey Effort..... | A-7 |
| Map 8: | Significant Fauna Database Search Results..... | A-8 |
| Map 9: | Fauna Habitat and Significant Fauna Records..... | A-9 |

Appendices

| | |
|-------------------|---|
| Appendix A | Maps |
| Appendix B | Literature Review Summary |
| Appendix C | Licences and Permits |
| Appendix D | Fauna Desktop Assessment Results |
| Appendix E | Fauna Site Sheets |
| Appendix F | Significant Fauna Recorded During the Survey |



Acronyms and Abbreviations

| | |
|--------------------|--|
| °C | Degree Celsius |
| BC Act | <i>Biodiversity Conservation Act 2016</i> |
| BoM | Bureau of Meteorology |
| CR | Critically Endangered |
| DAWE | Department of Agriculture Water and Environment |
| DBCA | Department of Biodiversity, Conservation and Attractions |
| DCCEEW | Department of Climate Change, Energy, the Environment and Water |
| DEE | Department of the Environment and Energy |
| Desktop Study Area | The area that was studied during the desktop assessment encompassing the Survey Area and surrounds |
| DMIRS | Department of Mines, Industry Regulation and Safety |
| DoE | Department of the Environment |
| DPIRD | Department of Primary Industries and Regional Development |
| DSEWPaC | Department of Sustainability, Environment, Water, Population and Communities |
| DWER | Department of Water and Environmental Regulation |
| EIA | Environmental Impact Assessment |
| EN | Endangered |
| EP Act | <i>Environmental Protection Act 1986</i> |
| EPA | Environmental Protection Authority |
| EPBC Act | <i>Environment Protection Biodiversity and Conservation Act 1999</i> |
| ESA | Environmentally Sensitive Area |
| GIS | Geographic Information System |
| GPS | Global Positioning System |
| ha | Hectare |
| IBRA | Interim Biogeographic Regionalisation for Australia |
| IBSA | Index of Biodiversity Surveys for Assessments |
| km | Kilometres |
| Lat | Latitude |
| Long | Longitude |
| m | Metres |
| MA | Marine |
| MI | Migratory |
| mm | Millimetres |
| MMP | Mining and Management Program |
| MMPLG | Mining and Management Planning Liaison Group |
| MNES | Matters of National Environmental Significance |



| | |
|-------------|---------------------------------|
| OS | Other Specially Protected Fauna |
| P | Priority |
| PMST | Protected Matters Search Tool |
| Alcoa | Alcoa of Australia Ltd |
| SLR | SLR Consulting Australia |
| Survey Area | The area that was surveyed |
| T | Threatened |
| TEC | Threatened Ecological Community |
| VU | Vulnerable |
| WA | Western Australia |
| WAM | Western Australian Museum |



1.0 Introduction

1.1 The Project

Alcoa of Australia Ltd (Alcoa) commissioned SLR Consulting Australia Pty Ltd (SLR) to undertake a fauna desktop assessment and targeted Chuditch (*Dasyurus geoffroii*), Quokka (*Setonix brachyurus*), and Woylie (*Bettongia penicillata*) survey for the Willowdale Mine Larego region (the Project) as part of its Mining and Management Program (MMP) assessment by the Mining and Management Planning Liaison Group (MMPLG). The survey was undertaken within a defined area that consists of approximately 35 individual polygons (the Survey Area) that cover 451.8 hectares and is located approximately 15 km southeast of the Waroona townsite, in the Jarrah Forest bioregion of Western Australia (**Map 1**). All maps are provided in **Appendix A**.

1.2 Objective and Scope

The objective of the survey was to identify key fauna values targeting Chuditch, Quokka, and Woylie within the Survey Area as part of the MMP assessment process for the Project. The following scope of work was completed:

- A desktop assessment of the Willowdale Mine Larego region including an overview of fauna habitat, inventory of likely species present and background environmental information.
- A targeted survey for Chuditch, Quokka, and Woylie.
- A technical biological report.



2.0 Background

2.1 Statutory and Regulatory Framework

Western Australian fauna is governed by the following legislative measures:

- *Environment Protection and Biodiversity Conservation Act 1999* (Cth) (EPBC Act). (Commonwealth of Australia, 1999).
- *Biodiversity Conservation Act 2016* (WA) (BC Act) (Government of Western Australia, 2016).
- *Environmental Protection Act 1986* (WA) (EP Act) (Government of Western Australia, 1986).

In addition to these legislative measures, the following non-legislative lists are considered on a case-by-case basis:

- WA Department of Biodiversity Conservation and Attractions (DBCA) Priority lists for fauna, flora, and ecological communities.
- Recognition of locally significant populations by DBCA.

The EIA process is supported by guidance documents published by the Environmental Protection Authority (EPA), DBCA and the Department of Climate Change, Energy, the Environment and Water (DCCEE).

Western Australia

- *Technical Guidance – Terrestrial vertebrate fauna surveys for environmental impact assessment* (EPA, 2020).
- *Environmental Factor Guideline – Terrestrial Fauna* (EPA, 2016).

Commonwealth

- *Matters of National Environmental Significance – Significant Impact Guidelines 1.1* (DoE, 2013).
- *Survey guidelines for Australia’s threatened mammals* (DSEWPoC, 2011).

2.2 Existing Environment

2.2.1 Climate


The closest long-term Bureau of Meteorology (BoM) weather station with a complete dataset is Dwellingup Weather Station (Station 9538), located approximately 17.1 km southwest of the Survey Area.

Climate statistics were calculated using data from the most current climate normal, which is defined as a 30-year interval where possible. A climate normal is a period long enough to include year-to-year variations while avoiding the influence of longer-term changes in climate (BoM, 2007).

The long-term (1957 to 2024) mean minimum temperature for Dwellingup Weather Station ranges from 5.5°C (August) to 14.6°C (February) and the long-term mean maximum temperature ranges from 15.1°C (July) to 39.8°C (January) (BoM, 2024) (**Graph 1**).



The Dwellingup Weather Station recorded 1009 mm of rainfall in the 12 months prior to the survey (October 2022 to September 2023), which is 209.3 mm below the long-term (1934 to 2023) average of 1218.3 mm (BoM, 2024). In the three months prior to the survey (July 2023 to September 2023), 459.4 mm of rainfall was recorded, which is 97.5 mm below the long-term average of 556.9 mm for the same period (BoM, 2024).



Graph 1: Climate summary of Dwellingup Weather Station

2.2.2 Interim Biogeographic Regionalisation of Australia

The Interim Biogeographic Regionalisation of Australia (IBRA) divides Australia into 89 bioregions based on major biological, geographical, and geological attributes. These bioregions are subdivided into 419 subregions as part of a refinement of the IBRA framework (DEE, 2016). The Survey Area occurs within the Jarrah Forest bioregion and the Northern Jarrah Forest (JAF01) subregion.

The Northern Jarrah Forest subregion overlays Archaean granite and metamorphic rocks with an average elevation of 300 m, capped by extensive lateritic duricrust and interrupted by later drainage and dissected by occasional granite hills. In the east the laterite becomes greatly separated until it compresses isolated remnants. Vegetation comprises of Jarrah-Marri Forest in the west with Bullich and Blackbutt in the valleys grading to Wandoo and Marri woodlands in the east with Powder bark on breakaways. A common understorey of heath is found on granite rocks in the north and east forests and woodlands. The lower slopes, where rapid changes in the site conditions occur, support most of the diversity in the communities. The subregional area is 2,255,904 ha (Williams and Mitchell, 2001).

2.2.3 Soil Landscape Mapping

Soil landscape mapping of Western Australia consists of a compilation of various surveys at different scales varying between 1:20,000 and 1:3,000,000 (DPIRD, 2022). The mapping comprises a nested hierarchy of levels, with each level a subdivision of the preceding level. Soil landscape mapping provides an indication of potential vegetation types and fauna



habitats within the Survey Area and has been described below to the highest level of detail available for the Survey Area.

The Survey Area occurs within one soil landscape system, the Darling Plateau System (255Dp) comprising of lateritic plateau, duplex sandy gravels, loamy gravels, wet soils, Jarrah-Marri-Wandoo forest and woodland.

The Survey Area occurs across four soil landscape subsystems and two soil landscape phases (**Table 1**; **Map 2**).

Table 1: Soil landscape subsystems and phases within the Survey Area

| Subsystem or phase | | Description | Area (ha) and percentage within Survey Area |
|---|-----------|--|---|
| Name | Code | | |
| Dwellingup Subsystem | 255DpDW | Divides, lower to upper slopes and hillcrests. Duplex sandy gravels and loamy gravels with minor areas of shallow gravels, deep sandy gravels, yellow deep sands, and yellow and pale deep sands, often gravelly | 364.0 ha (80.6%) |
| Hester Subsystem | 255DpHR | Ridges and hill crests on laterite and gneiss, relief 5-40 m, slopes 5-15%. Soils are sandy gravels, loamy gravels, and loamy earths | 3.9 ha (0.9%) |
| Mornington Hill Subsystem | 255DpMH | Low hills on laterite overlying granite, relief 40-80 m, slope 5-20%. Soils are sandy and loamy gravels with some deep sands and loamy earths | 23.6 ha (5.2%) |
| Yarragil DpYGh phase | 255DpYGh | Very gentle to moderately inclined (<20%) concave valley sideslopes. Moderately well drained yellow duplex soils and yellow and brown massive earths | 20.2 ha (4.5%) |
| Yarragil subsystem | 255DpYG | Shallow, narrow, upper valleys of the deeply dissected Murray, Bindoon, and Helena units. Alluvial, clay and loam soils, moderately well drained, often gravelly, with some sands and loams. Salt prone. Woodland of <i>E. wandoo</i> , <i>E. accedens</i> | 39.0 ha (8.6%) |
| Yarragil Swamp phase | 255DpYGsw | Level to very gently inclined valley floors. Swampy river flats and terraces in granitic rocks; loamy and sandy duplex, wet soils, non-cracking clays, and loams | 1.0 ha (0.2%) |
| Total area and percentage within Survey Area | | | 451.8 ha (100%) |

2.2.4 Hydrography

A total of two hydrographic features intersect the Survey Area, and a total of nine hydrographical features occur within a 5 km radius of the Survey Area are described in **Table 2** and shown in **Map 3** (DWER, 2018).

Table 2: Hydrographic features in the vicinity of the Survey Area

| Hydrographical feature | Description | |
|------------------------|--------------------|----------------|
| Bancell Brook | Significant Stream | 4 km west |
| Clarke Brooke | Significant Stream | 5 km southwest |



| Hydrographical feature | Description | |
|----------------------------------|------------------------|--|
| Harvey River | Mainstream | Intersects two polygons in the west of the Survey Area |
| Logue Brook | Minor River | 3.0 km southwest |
| Logue Brook Dam | Water Supply Reservoir | 2.9 km west |
| Samson Brook Dam – Lake Kabbanup | Water Supply Reservoir | 3.4 km north |
| Samson Brook | Signification stream | 0.4 km north |
| Unnamed Significant Stream | Significant stream | 3.7 km northwest |
| Unnamed Tributary | Major tributary | Intersects the eastern polygon of the Survey Area |
| Unnamed Tributary | Minor tributary | 5.5 km west |
| Unnamed Tributary | Minor tributary | 4.0 km west |

2.2.5 Pre-European Vegetation

The major source of data for pre-European vegetation mapping in Western Australia is the published and unpublished mapping of J. S. Beard at 1:250,000 scale. These vegetation types were later refined by Shepherd, Beeston, and Hopkins (2002), resulting in 819 Vegetation Association-level units, and a subsequent reclassification resulted in the creation of over 2,175 finer-scale System Associations (Beard *et al.*, 2013). The Survey Area occurs within one System Association, WEST DARLING_3, which comprises Jarrah (*Eucalyptus marginata*) and Marri (*Corymbia calophylla*) (**Map 4**).

2.2.6 Conservation Areas

Conservation areas consist of areas protected for the purpose of conservation, including but not limited to National Parks, Nature Reserves, Conservation Parks, and Regional Parks. The Survey Area occurs within the Dwellingup State Forest which is vested under Conservation Commission of WA (DBCA, 2023a, 2023b) and shown in **Map 5**.

2.2.7 Environmentally Sensitive Areas

Environmentally Sensitive Areas (ESAs) are declared by the Department of Water and Environmental Regulation (DWER) to prevent the degradation of important environmental values such as Threatened flora, Threatened Ecological Communities (TECs), or significant wetlands. The Survey Area does not intersect any mapped ESAs (DWER, 2021). There are nine Swan Coastal Plain wetland ESAs within a 5 km radius of the Survey Area. ESAs are shown in **Map 5**.

2.2.8 Land Use

No pastoral leases apply to the Survey Area. The Survey Area is under one mining tenement, Alcoa of Australia Ltd (AML7000001) (DMIRS, 2024). Three pending exploration licences also intersect the Survey Area:

- Eastern Goldfields Exploration Pty Ltd E7002338.
- G E Resources Pty Ltd E7005390.
- G E Resources Pty Ltd E7005543.



3.0 Methods

The surveys documented in this report were undertaken in accordance with relevant EPA and DCCEEW guidelines (see **Section 2.1**).

3.1 Desktop Assessment

3.1.1 Literature Review

Background information on the Survey Area and surrounds (the Desktop Study Area) was compiled prior to the field survey. The literature review considered a selection of relevant reports detailing assessments undertaken in the region that were either publicly available from sources such as the Index of Biodiversity Surveys for Assessments (IBSA) website, the EPA Consultation Hub, or internet searches, or provided by Alcoa. These reports are listed below, summarised in **Appendix B**, and their locations in relation to the current Survey Area, as represented by their nearest point, are shown in **Map 6**:

- *Fauna Assessment Pinjarra - Williams Road (SLK 14 – 67)* (Harewood, 2017), 14 km north of the Survey Area.
- *Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Holyoake* (GHD, 2021a), 18 km north of the Survey Area.
- *Black Cockatoo Habitat Assessment – Pinjarra Racecourse* (Harewood, 2019), 30 km north of the Survey Area.
- *Chuditch Pre-mine Survey, Myara* (Way, 2013), 43 km north of the Survey Area.
- *Vertebrate Fauna Survey of Planned Mining Areas at Alcoa’s Willowdale Larego Mining Region* (Environmental Management and Research Consultants, 2006), 43 km north of the Survey Area.
- *Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Myara North* (GHD, 2021b), 54 km north of the Survey Area.
- *Report for Rail Reserves in the Shire of Serpentine Jarrahdale - Spring Flora and Vegetation Survey and Fauna and Habitat Assessment* (GHD, 2012), 60 km north of the Survey Area.

3.1.2 Database Searches

Database searches were undertaken to compile a list of fauna known to occur in the Desktop Study Area and identify significant fauna with potential to occur within the Survey Area (**Table 3**).

Table 3: Database search details

| Database name | Date received | Search target | Buffer around the Survey Area |
|---|---------------|---|-------------------------------|
| Threatened and Priority Fauna database search (DBCA, 2023c) | 08 June 2023 | Threatened and Priority fauna | 20 km |
| Protected Matters Search Tool (PMST) (DCCEEW, 2023) | 14 July 2023 | Threatened flora, fauna, and ecological communities | 20 km |



| Database name | Date received | Search target | Buffer around the Survey Area |
|-------------------------|---------------|---------------|-------------------------------|
| NatureMap (DBCA, 2023d) | 07 June 2023 | Fauna | 20 km |

3.1.3 Likelihood of Occurrence

Significant fauna taxa identified during the desktop assessment were assessed to determine the likelihood of their occurrence within the Survey Area before and after the field survey. The assessment used the likelihood of occurrence criteria presented in **Table 4**.

Taxa listed as Marine only under the EPBC Act were not considered to be significant taxa because the Marine listing does not constitute MNES under the EPBC Act. Additionally, erroneous records (i.e. records that occur well outside a taxon's known distribution) were excluded from consideration. Only taxa that have been recorded within the Survey Area or were assessed as having a high or medium likelihood of occurrence are discussed in detail.

Table 4: Likelihood of occurrence criteria

| Rank | Criteria |
|----------------------------|--|
| Recorded | The taxon was recorded within the Survey Area during the current survey. |
| Previously Recorded | The taxon has been previously recorded within the Survey Area according to database search or literature review results. |
| High (Likely to occur) | The taxon has been recorded within the Desktop Study Area more than once in the last 15 years and suitable habitat is present within the Survey Area. |
| Medium (May occur) | There are existing records of the taxon within the Desktop Study Area, however the taxon has not been recorded within the Desktop Study Area in the last 15 years or suitable habitat within the Survey Area is marginal or limited in extent. |
| Low (Unlikely to occur) | Suitable habitat is not present within the Survey Area, or the taxon is very infrequently recorded in the locality despite reasonable previous search effort, or the taxon is believed to be extinct or locally extinct. |

3.2 Field Survey

3.2.1 Survey Timing

The field surveys were undertaken across two field trips as shown in **Table 5**.

Table 5: Survey timing

| Survey trip | Tasks completed | Dates | Person field days |
|-------------|--|--|-------------------|
| 1 | Active searches, deployment of camera traps, habitat assessment, spotlighting. | 13 th October – 16 th October 2023 | 16 |
| 2 | Collection of camera traps | 7 th November – 8 th November 2023 | 4 |



3.2.2 Field Personnel and Licences

Details of field personnel, including their level of experience and role for each field trip are detailed in **Table 6**. Fieldwork was completed under Fauna Taking (Biological Assessment) License – Regulation 27 (08-005511-1) (**Appendix C**).

Table 6: Field personnel

| Personnel | Experience | Role | Trips |
|-----------------|------------|---------------------|---------|
| Dr Michael Lohr | 11 years | Principal Zoologist | 1 |
| Evan Webb | 7 years | Associate Zoologist | 1 and 2 |
| Poppy Walker | 5 years | Senior Zoologist | 2 |
| Lewis Berry | 2 years | Ecologist | 1 |
| Datta Li | 2 years | Zoologist | 1 |

3.2.3 Habitat Assessment and Mapping

Habitat assessments were undertaken in representative areas of fauna habitat within the Survey Area to record habitat values. A habitat assessment was recorded at each camera trap site, and two additional standalone habitat assessments were recorded. Habitat assessment locations are shown in **Map 7**.

The following information was collected at each habitat assessment locations using a GPS-enabled handheld device:

- Site photo.
- Landform.
- Soil type and colour.
- Rock types, surface stone cover, and size classes.
- Key habitat and microhabitat features including leaf litter, logs, burrows, rocky outcrops, rock crevices, hollows, and water sources.
- Habitat quality, fire history, and evidence of disturbance.
- General description of vegetation structure.

Fauna habitat mapping was based on a combination of field observations, habitat assessment data, aerial imagery, and vegetation mapping. Polygons were digitised using GIS software.

3.2.4 Camera Traps

A total of 45 motion sensitive camera traps were deployed during the field survey. Each camera trap was deployed for a minimum of 23 nights and total survey effort consisted of 1,119 trap nights. Cameras were placed in areas of suitable habitat for (Chuditch, Quokka, and Woylie) and non-target taxa (Wambenger Brush-tailed Phascogale, Quenda, and Western Brush Wallaby) and were baited with universal bait (rolled oats, peanut butter, and sardines) and chopped apples. Camera trap locations are shown in **Map 7**.

3.2.5 Spotlighting

Spotlighting was undertaken on two nights during the field survey. Spotlighting commenced approximately one hour after sunset and consisted of traversing areas of suitable habitat, generally near stream crossings, on foot for a duration of two hours. A total of 16 person



hours was completed during the field survey. Spotlighting was also undertaken opportunistically from the vehicle while travelling to and from the spotlighting location. Spotlighting was undertaken at the following locations (**Map 7**):

- 5890-CAM-02-EW
- 5890-CAM-20-EW
- 5890-CAM-21-EW
- 5890-CAM-44-EW

3.2.6 Opportunistic Observations

Opportunistic observations of fauna were recorded throughout the Survey Area, including primary evidence (direct sightings, calls) and secondary evidence (tracks, scats, diggings, remains).

3.2.7 Active Searches

Untimed active searches were undertaken at each camera trap site. These searches included flipping rocks and logs and raking leaf litter in search of direct sightings, scats, tracks, and other traces of fauna, with a particular focus on Chuditch (scats, dens in hollow logs or burrows), Quokka (scats, runnels in sedges), Woylie (scats, diggings, nests), Numbat (scats, diggings), and Western Ringtail Possums (dreys, scats).

3.3 Limitations

Limitations and constraints of the fauna survey are detailed below in **Table 7**.

Table 7: Limitations and constraints associated with the survey

| Variable | Degree of limitation | Potential constraints on survey outcomes |
|---|----------------------|---|
| Availability of data and information | No | Sufficient data and information, including regional and local contextual information, was available to complete the scope of the survey. |
| Competency and experience of the survey team | No | The survey was undertaken by a team with extensive experience undertaking similar scopes within the bioregion. <ul style="list-style-type: none"> • Principal Zoologist Dr Michael Lohr – 11 years' experience. • Associate Zoologist Evan Webb – 7 years' experience. • Senior Zoologist Poppy Walker – 5 years' experience. • Zoologist Datta Li – 2 years' experience. • Ecologist Lewis Berry – 2 years' experience. |
| The proportion of fauna identified, recorded, or collected | No | All targeted fauna taxa were able to be identified to species level with a high degree of confidence. |
| Scope of the survey | No | The scope of the survey was limited to terrestrial vertebrate fauna targeting threatened marsupials. No further exclusions were made within this group. |
| Adequacy of the survey intensity and proportion of survey achieved | No | Sufficient time and effort were allocated to the survey given the size and complexity of the Survey Area and the expected level of survey intensity. |



| Variable | Degree of limitation | Potential constraints on survey outcomes |
|---|----------------------|---|
| Access problems | No | The Survey Area was sufficiently accessed by vehicle and on foot. |
| Timing and season | No | Targeted Chuditch, Quokka, and Woylie surveys can be undertaken at any time of year. |
| Disturbance that may have affected the results of survey | No | Areas of disturbance associated with mining activities and bushfires were present within the Survey Area but were not a limitation on the results of the survey. |
| Problems with data and analysis, including sampling biases | No | Survey effort for significant fauna taxa was concentrated in preferred habitats. This may introduce bias where the use of non-preferred habitat is underrepresented, however, this is not considered a limitation on the survey outcomes. |



4.0 Results

4.1 Desktop Assessment

The database searches and literature review identified 189 terrestrial vertebrate fauna taxa occurring within the Desktop Study Area, of which 33 are significant (Threatened or Priority):

- 126 birds, of which 19 are significant.
- 28 mammals, of which 10 are significant.
- 25 reptiles, of which 4 are significant.
- 10 amphibians, of which none are significant.



Key findings of the literature review are summarised in **Appendix B**, a complete list of fauna taxa recorded within the Desktop Study Area is presented in **Appendix D**, and DBCA database search results are displayed in **Map 8**.

4.2 Fauna Habitat

Two fauna habitats (excluding cleared areas) were identified and mapped within the Survey Area. Fauna habitats are presented in **Map 9**, described below in **Table 8**, and site sheets for each habitat assessment are provided in **Appendix E**. Small discrepancies in fauna habitat extents (i.e., not adding up to the exact area extent of the Survey Area) are due to rounding.



Table 8: Fauna habitats recorded within the Survey Area

| Fauna habitat | Total area, percentage of Survey Area | Habitat description | Representative photograph |
|---------------------|---------------------------------------|--|--|
| Jarrah-Marri Forest | 439.2 ha, 97.2% | <p>Undulating plains and hills over clay loam soils and laterite rock. Vegetation consists of Jarrah (<i>Eucalyptus marginata</i>) and Marri (<i>Corymbia calophylla</i>) open forest with patchy <i>Allocasuarina Fraseriana</i> over a mixed midstorey typically dominated by <i>Bossiaea aquifolium</i>, <i>Xanthorrhoea preissi</i>, and <i>Macrozamia fraseri</i>, over sedges and bracken. Disturbances consisted primarily of historic logging and bushfire. Microhabitats present within this habitat include hollows in trees and logs, leaf litter and woody debris.</p> <p>This habitat is core habitat Chuditch and supporting habitat for Woylie and Quokka. The canopy is core habitat for black cockatoos.</p> |  |
| Drainage Line | 12.7 ha, 2.8% | <p>Areas of drainage over clay loam soils, often with thin channels of surface water present. Vegetation consists of a closed mosaic upperstorey of Bullich (<i>E. megacarpa</i>), Blackbutt (<i>E. patens</i>), and Flooded Gum (<i>E. rudis</i>), forest over a closed mixed shrubland of <i>Melaleuca</i> sp., <i>B. aquifolium</i>, and other <i>Myrtaceae</i> sp. and <i>Acacia</i> sp., over tall sedgeland.</p> <p>This habitat typically had low levels of disturbance. Microhabitats present within this habitat include tree and log hollows, woody debris, peeling bark and leaf litter.</p> <p>Drainage Line habitat is core habitat Chuditch, Woylie, and Quokka and the canopy constitutes core habitat for black cockatoos. This habitat is analogous to Blackbutt, Bullich, and Flooded Gum fauna habitats recorded by GHD (2021b, 2021a).</p> |  |
| Total | 451.8 ha, 100% | | |



4.3 Recorded Fauna

The basic terrestrial vertebrate fauna survey recorded 30 fauna taxa from 19 families, comprising three amphibian taxa, 12 bird taxa, 10 mammal taxa, and five reptile taxa. Of the taxa recorded, five were significant (see **Section 4.4**). Three introduced taxa were recorded, the Cat (*Felis catus*), Black Rat (*Rattus rattus*), and Pig (*Sus scrofa*).

An inventory of fauna recorded during the field survey is provided below in **Table 9**.



Table 9: Fauna recorded within the Survey Area

| Family | Scientific Name | Common Name | Conservation Status | | Call | Camera Trap | Capture | Feather | Foraging evidence | Scat | Sighting | Sighting - overhead | Total |
|-------------------|--|----------------------------------|---------------------|---------------|------|-------------|---------|---------|-------------------|------|----------|---------------------|-------|
| | | | State | Common wealth | | | | | | | | | |
| Amphibians | | | | | | | | | | | | | |
| Limnodynastidae | <i>Heleioporus inornatus</i> | Whooping Frog | - | - | | | | | | | 1 | | 1 |
| Myobatrachidae | <i>Crinia georgiana</i> | Quacking Frog | - | - | 7 | | | | | | 1 | | 8 |
| Myobatrachidae | <i>Crinia glauerti</i> | Clicking Frog | - | - | 10 | | | | | | 1 | | 11 |
| Birds | | | | | | | | | | | | | |
| Acanthizidae | <i>Gerygone fusca</i> | Western Gerygone | - | - | 1 | | | | | | | | 1 |
| Acanthizidae | <i>Sericornis maculatus</i> | Spotted Scrubwren | - | - | | 7 | | | | | | | 7 |
| Cacatuidae | <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | | | | 3 | 2 | | 12 | 3 | 20 |
| Cacatuidae | <i>Zanda baudinii</i> | Baudin's Cockatoo | EN | EN | | | | | 1 | | | | 1 |
| Columbidae | <i>Phaps elegans</i> | Brush Bronzewing | - | - | | 3 | | | | | | | 3 |
| Maluridae | <i>Malurus elegans</i> | Red-winged Fairywren | - | - | | 26 | | | | | | | 26 |
| Maluridae | <i>Malurus splendens</i> | Splendid Fairywren | - | - | | 1 | | | | | | | 1 |
| Pachycephalidae | <i>Colluricincla harmonica</i> | Grey Shrikethrush | - | - | 1 | | | | | | | | 1 |
| Petroicidae | <i>Eopsaltria griseogularis</i> | Western Yellow Robin | - | - | | 1 | | | | | | | 1 |
| Petroicidae | <i>Quoyornis georgianus</i> | White-breasted Robin | - | - | | 5 | | | | | | | 5 |
| Rhipiduridae | <i>Rhipidura albiscapa</i> | Grey Fantail | - | - | | 2 | | | | | | | 2 |
| Strigidae | <i>Ninox boobook</i> | Boobook Owl | - | MA | 1 | | | | | | | | 1 |
| Mammals | | | | | | | | | | | | | |
| Dasyuridae | <i>Antechinus flavipes leucogaster</i> | Mardo | - | - | | 78 | | | | | | | 78 |
| Dasyuridae | <i>Dasyurus geoffroii</i> | Chuditch | VU | VU | | | | | | 1 | | | 1 |
| Felidae | * <i>Felis catus</i> | Cat | - | - | | 1 | | | | | | | 1 |



| Family | Scientific Name | Common Name | Conservation Status | | Call | Camera Trap | Capture | Feather | Foraging evidence | Scat | Sighting | Sighting - overhead | Total |
|-----------------|---|-----------------------------|---------------------|---------------|------|-------------|---------|---------|-------------------|------|----------|---------------------|-------|
| | | | State | Common wealth | | | | | | | | | |
| Macropodidae | <i>Macropus fuliginosus melanops</i> | Western Grey Kangaroo | - | - | | 10 | | | | | | | 10 |
| Macropodidae | <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | - | | 26 | | | | | 3 | | 29 |
| Muridae | <i>Hydromys chrysogaster</i> | Rakali | P4 | - | | 1 | | | | | | | 1 |
| Muridae | * <i>Rattus rattus</i> | Black Rat | - | - | | 70 | | | | | | | 70 |
| Phalangeridae | <i>Trichosurus vulpecula hypoleucus</i> | Common Brushtail Possum | - | - | | 1 | | | | | | | 1 |
| Suidae | * <i>Sus scrofa</i> | Pig | - | - | | 16 | | | | | 1 | | 17 |
| Tachyglossidae | <i>Tachyglossus aculeatus acanthion</i> | Short-beaked Echidna | - | - | | 17 | | | | | | | 17 |
| Reptiles | | | | | | | | | | | | | |
| Scincidae | <i>Egernia napoleonis</i> | South-western Crevice-skink | - | - | | 93 | 1 | | | | 2 | | 96 |
| Scincidae | <i>Lissolepis luctuosa</i> | Western Swamp Skink | - | - | | 1 | | | | | 1 | | 2 |
| Scincidae | <i>Morethia obscura</i> | Shrubland Morethia Skink | - | - | | | 1 | | | | | | 1 |
| Scincidae | <i>Tiliqua rugosa</i> | Bobtail | - | - | | 51 | | | | | | | 51 |
| Varanidae | <i>Varanus gouldii</i> | Bungarra | - | - | | 1 | | | | | 1 | | 2 |

* = introduced species



4.4 Significant Fauna

Five significant fauna taxa were recorded within the Survey Area during the field survey. These are listed below, and locations are presented in **Map 9** and **Appendix F**:

- Baudin's Cockatoo (*Zanda baudinii*) – EN (BC Act; EPBC Act), recorded once within Jarrah-Marri Forest habitat.
- Forest Red-tailed Black Cockatoo (*Calyptorhynchus banksii naso*) – VU (BC Act; EPBC Act), recorded 20 times within Jarrah-Marri Forest and Drainage Line habitats.
- Chuditch (*Dasyurus geoffroyi*) – VU (BC Act; EPBC Act), recorded once within Jarrah-Marri Forest habitat (**Figure 1**).
- Rakali (*Hydromys chrysogaster*) – P4 (DBCA), recorded once within Drainage Line habitat (**Figure 2**).
- Western Brush Wallaby (*Notamacropus irma*) – P4 (DBCA), recorded 29 times within Jarrah-Marri Forest and Drainage Line habitats (**Figure 3**).

Seven significant fauna taxa were assessed as having a high likelihood of occurring within the Survey Area:

- Woylie (*Bettongia penicillata ogilbyi*) – CR (BC Act); EN (EPBC Act)
- Carnaby's Cockatoo (*Zanda latirostris*) – EN (BC Act; EPBC Act).
- Quokka (*Setonix brachyurus*) – VU (BC Act; EPBC Act)
- Wambenger Brush-tailed Phascogale (*Phascogale tapoatafa wambenger*) – CD (BC Act).
- Quenda (*Isoodon fusciventer*) – P4 (DBCA).
- Western False Pipistrelle (*Falsistrellus mackenziei*) – P4 (DBCA).
- Dell's Ctenotus (*Ctenotus delli*) – P4 (DBCA).

Six significant fauna taxa were assessed as having a medium likelihood of occurring within the Survey Area:

- Western Ringtail Possum (*Pseudocheirus occidentalis*) – CR (BC Act; EPBC Act).
- Australasian Bittern (*Botaurus poiciloptilus*) – EN (BC Act).
- Malleefowl (*Leipoa ocellata*) – VU (BC Act; EPBC Act).
- Peregrine Falcon (*Falco peregrinus*) – OS (BC Act).
- Woma Python (*Aspidites ramsayi*) – P1 (DBCA).
- Southern Death Adder (*Acanthophis antarcticus*) – P4 (DBCA).

The complete results of the significant fauna likelihood of occurrence assessment including justification for the assessment outcome for each taxon is provided in **Table 10**.





Figure 1: Chuditch scat recorded within Jarrah-Marri Forest (-32.9758445, 116.0860574)



Figure 2: Rakali recorded by camera trap 5890-CAM-07-EW within Drainage Line habitat (-32.9941287, 116.0524062).





Figure 3: Western Brush Wallaby recorded by camera trap 5870-CAM-12-LB within Jarrah-Marri Forest (-32.97344506, 116.0826827).



Table 10: Significant fauna likelihood of occurrence within the Survey Area

| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|--------------|---|---------------------|--------------|---|--|--|
| | | State | Commonwealth | | | |
| Birds | | | | | | |
| Acanthizidae | <i>Aphelocephala leucopsis</i> Southern Whiteface | - | VU | Sparse open forest/woodland; inland scrubs e.g. mallee, mulga cypress pine; saltbush, dead trees, stumps, black box/lignum flats (Pizzey and Knight, 2001). | No nearby records identified. | Low No nearby records and limited suitable habitat within the Survey Area. |
| Accipitridae | <i>Elanus scriptus</i> Letter-winged Kite | P4 | - | Open country and grasslands of arid and semi-arid interior (Pizzey and Knight, 2001). | No DBCA records, one record from literature 60 km north of the Survey Area in 2011 (GHD, 2012). | Low No suitable habitat within the Survey Area. |
| Anatidae | <i>Oxyura australis</i> Blue-billed Duck | P4 | - | Densely vegetated freshwater lakes, swamps, dams (Morcombe, 2003). | Five DBCA records within 20 km of the Survey Area. Including 15.8 km southwest of the Survey Area in 2013 (DBCA, 2023c). | Low No suitable habitat within the Survey Area. |
| Ardeidae | <i>Botaurus poiciloptilus</i> Australasian Bittern | EN | EN | Freshwater wetlands with heavy vegetation, shrubbery, reedbeds, sedges (Morcombe, 2003). | Two DBCA records within 20 km of the Survey Area, including 5.8 km north in 2017 (DBCA, 2023c). | Medium One record 5.8 km from the Survey Area, limited suitable habitat within the Survey Area. |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|------------------|---|---------------------|--------------|---|--|---|
| | | State | Commonwealth | | | |
| Atrichornithidae | <i>Atrichornis clamosus</i> Noisy Scrubbird | EN | EN | Dense heaths, rushes, tall sedges, under stunted trees in coastal hills, coastal swamps (Morcombe, 2003). | Forty-four DBCA records within 20 km of the Survey Area, including eight records within the 1 km of the Survey Area as recent as 2007, many of which are translocated individuals (DBCA, 2023c). | Low No nearby records within the last 15 years and limited suitable habitat within the Survey Area. Individuals were translocated to the northern Jarrah forest in 1997 – 2003 but are now considered locally extinct (GHD, 2021b, 2021a). |
| Cacatuidae | <i>Calyptorhynchus banksii naso</i> Forest Red-tailed Black Cockatoo | VU | VU | Tall eucalypt forest, woodland, feeds on seeds of large-fruited eucalypts (Menkhorst <i>et al.</i> , 2019). | Eighty-five DBCA records within 20 km of the Survey Area, including two records within 1 km of the Survey Area as recent as 2018 (DBCA, 2023c). | Recorded Recorded during the current survey. |
| Cacatuidae | <i>Zanda baudinii</i> Baudin's Cockatoo | EN | EN | Forests, farm trees; feeds on Marri and wood-boring insects (Morcombe, 2003). | Fifty DBCA records within 20 km of the Survey Area, including two records within 1 km of the Survey Area as recent as 2002 (DBCA, 2023c). | Recorded Recorded during the current survey. |
| Cacatuidae | <i>Zanda latirostris</i> Carnaby's Cockatoo | EN | EN | Forests, woodlands, heathlands, farms; feeds on Banksias and Hakeas, pine plantations (Morcombe, 2003). | Forty-eight DBCA records within 20 km of the Survey Area, including three records within 1 km of the Survey Area as recent as 2007 (DBCA, 2023c). | High Nearby records, and suitable habitat within the Survey Area. |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|--------------|--|---------------------|--------------|--|---|--|
| | | State | Commonwealth | | | |
| Charadriidae | <i>Charadrius leschenaultia</i> Greater Sand Plover | VU | VU, MI, MA | Tidal flats, beaches . | No nearby records identified. | Low |
| | | | | | | No nearby records, and no suitable habitat within the Survey Area. |
| Charadriidae | <i>Thinornis cucullatus</i> Hooded Dotterel | P4 | MA | Beaches, margins of inland salt lakes (Menkhorst <i>et al.</i> , 2019). | No DBCA or literature records, one Naturemap record within 20 km of the Survey Area (DBCA, 2023d). | Low |
| | | | | | | One NatureMap record, and no suitable habitat within the Survey Area. |
| Falconidae | <i>Falco hypoleucos</i> Grey Falcon | VU | VU | Open plains with treed watercourses in arid inland (Menkhorst <i>et al.</i> , 2019). | No nearby records identified. | Low |
| | | | | | | No nearby records, and no suitable habitat within the Survey Area. |
| Falconidae | <i>Falco peregrinus</i> Peregrine Falcon | OS | - | Most environments with suitable nest sites: cliff faces preferred, including man-made ones, commonly uses stick nests built by other species (Menkhorst <i>et al.</i> , 2019). | Two DBCA records within 20 km of the Survey Area, including 15 km south of the Survey Area in 2007 (DBCA, 2023c). | Medium |
| | | | | | | Records <15 km from the Survey Area and may use the Survey Area for hunting. |
| Laridae | <i>Sternula nereis</i> Fairy Tern | VU | VU | Coastal, bays, inlets, beaches, salt ponds and lakes (Morcombe, 2003). | No nearby records identified. | Low |
| | | | | | | No suitable habitat within the Survey Area. |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|---------------|---|---------------------|--------------|--|---|--|
| | | State | Commonwealth | | | |
| Megapodiidae | <i>Leipoa ocellata</i> Malleefowl | VU | VU | Unburned mallee and woodland with abundant litter and low scrub (Morcombe, 2003). | Two DBCA records within 20 km of the Survey Area, including 17.6 km north of the Survey Area in 2014 (DBCA, 2023c). | <p style="text-align: center;">Medium</p> <p>Outside known distribution, however, taxon has been sighted near the Survey Area. Furthermore, individuals were translocated to the nearby Dryandra Woodland, 79 km east-northeast of the Survey Area, within the last five years (Traill, 2019).</p> |
| Rostratulidae | <i>Rostratula australis</i> Australian Painted Snipe | EN | EN, MA | Well vegetated surrounds and shallows of wetlands (Morcombe, 2003). | No nearby records identified. | <p style="text-align: center;">Low</p> <p>No nearby records, and no suitable habitat within the Survey Area.</p> |
| Scolopacidae | <i>Calidris canutus</i> Red Knot | EN | EN, MI, MA | Coastal, extensive firm tidal flats (Menkhorst <i>et al.</i> , 2019). | No nearby records identified. | <p style="text-align: center;">Low</p> <p>No nearby records, and no suitable habitat within the Survey Area.</p> |
| Scolopacidae | <i>Calidris ferruginea</i> Curlew Sandpiper | CR | - | Inter-tidal mudflats of estuaries, lagoons, mangrove channels, dams, floodwaters, flooded saltbush surrounds of inland lakes (Morcombe, 2003). | No nearby records identified from the database searches or literature. | <p style="text-align: center;">Low</p> <p>No nearby records, and no suitable habitat within the Survey Area.</p> |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|----------------|--|---------------------|--------------|---|---|--|
| | | State | Commonwealth | | | |
| Scolopacidae | <i>Numenius madagascariensis</i> Eastern Curlew | CR | CR, MI, MA | Coastal, tidal flats (Menkhorst <i>et al.</i> , 2019). | No nearby records identified from the database searches or literature. | Low |
| | | | | | | No nearby records, and no suitable habitat within the Survey Area. |
| Tytonidae | <i>Tyto novaehollandiae</i> Australian Masked Owl | P3 | - | Forests and woodlands with hollow bearing large old trees; roosts in tree hollows, foliage, and caves (Menkhorst <i>et al.</i> , 2019). | One DBCA record 13.6 km north of the Survey Area in 1971 (DBCA, 2023c). | Low |
| | | | | | | No recent nearby records, suitable habitat within the Survey Area. |
| Mammals | | | | | | |
| Dasyuridae | <i>Dasyurus geoffroii</i> Chuditch | VU | VU | Sclerophyll forest or drier woodland, heath, and mallee shrubland (Van Dyck and Strahan, 2008). | Eighty-eight DBCA records within 20 km of the Survey Area, and one record 1.7 km northeast from the Survey Area (DBCA, 2023c). | Recorded |
| | | | | | | Recorded during the current survey. |
| Dasyuridae | <i>Phascogale tapoatafa wambenger</i> Wambenger Brush-tailed Phascogale | CD | - | Mature rough-barked trees, large logs, dead standing trees (Van Dyck and Strahan, 2008). | Seventeen DBCA records within 20 km of the Survey Area, including one 13.1 km southwest of the Survey Area in 2022 (DBCA, 2023c). | High |
| | | | | | | Recorded <15 km and suitable habitat within the Survey Area. |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|---------------|--|---------------------|--------------|---|--|---|
| | | State | Commonwealth | | | |
| Macropodidae | <i>Setonix brachyurus</i> Quokka | VU | VU | Dense wet ground cover in forest or swampy flats (Van Dyck and Strahan, 2008). | Twenty-six DBCA records within 20 km of the Survey Area, including one record within 200 m of the Survey Area in 2016 (DBCA, 2023c). | High Recorded 200 m from the Survey Area, and suitable habitat within the Survey Area. |
| Macropodidae | <i>Notamacropus irma</i> Western Brush Wallaby | P4 | - | Open forest and woodland, open seasonally wet flats (Van Dyck and Strahan, 2008). | Nineteen DBCA records within 20 km of the Survey Area, including 14.8 km north in 2006 (DBCA, 2023c). | Recorded Recorded during the current survey. |
| Muridae | <i>Hydromys chrysogaster</i> Rakali | P4 | - | Permanent fresh to brackish water bodies (Van Dyck and Strahan, 2008). | Seven DBCA records within 20 km of the Survey Area, including one within 15.6 km north of the Survey Area in 2018 (DBCA, 2023c). | Recorded Recorded during the current survey. |
| Myrmecobiidae | <i>Myrmecobius fasciatus fasciatus</i> Numbat, Walpurti | EN | EN | Jarrah forests, Wandoo woodlands; requires hollow logs and branches for shelter and termites for food (Van Dyck and Strahan, 2008). | Two historic records from 1980 and 1954 within 20 km of the Survey Area. | Low No nearby recent records. |
| Peramelidae | <i>Isodon fusciventer</i> Quenda | P4 | - | Sandy soils with dense heathy vegetation (Van Dyck and Strahan, 2008). | Sixty-four DBCA records within 20 km of the Survey area, including records within 1.6 km of the Survey Area in 1995 (DBCA, 2023c). | High Nearby records, and suitable habitat within the Survey Area. |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|------------------|--|---------------------|--------------|---|---|---|
| | | State | Commonwealth | | | |
| Potoroidae | <i>Bettongia penicillata ogilbyi</i> Brush-tailed Bettong, Woylie | CR | EN | Areas dominated by <i>Gastrolobium</i> thickets (Van Dyck and Strahan, 2008). | Thirty-nine DBCA records within 20 km of the Survey Area, including 10 records within 500 m of the Survey Area as recent as 2004 (DBCA, 2023c). | High Recorded <500m from the Survey Area, and suitable habitat within the Survey Area. |
| Pseudocheiridae | <i>Pseudocheirus occidentalis</i> Western Ringtail Possum | CR | CR | <i>Agonis</i> forest and woodland, Tuart Forest with <i>Agonis</i> mid storey, and Jarrah, Marri and Wandoo woodlands (Van Dyck and Strahan, 2008). | Seventeen DBCA records within 18 km of the Survey Area, including 9.2 km north of the Survey Area in 2020 (DBCA, 2023c). | Medium Survey Area is outside of the taxon's distribution. Nearby records may be associated with translocations at the Lane Poole Reserve and other sites. |
| Vespertilionidae | <i>Falsistrellus mackenziei</i> Western False Pipistrelle | P4 | - | Mature Karri forests, wetter stands of Jarrah and Tuart, and woodlands on Swan Coastal Plain (Van Dyck and Strahan, 2008). | Four DBCA records within 15 km of the Survey Area, including two 8.8 km northwest of the Survey Area in 2013 (DBCA, 2023c). | High Records <9 km from the Survey Area, and suitable habitat within the Survey Area. |
| Reptiles | | | | | | |
| Elapidae | <i>Acanthophis antarcticus</i> Southern Death Adder | P3 | - | Habitats range from rainforest to shrublands and heath (Wilson and Swan, 2021). | One record 55 km north of the Survey Area in 2021 (GHD, 2021b). | Medium No nearby records, however, species is cryptic and suitable habitat present within the Survey Area. |



| Family | Scientific Name and Common Name | Conservation Status | | Habitat | Previous Records | Likelihood of Occurrence |
|------------|--|---------------------|---------------|--|--|--|
| | | State | Common wealth | | | |
| Pythonidae | <i>Aspidites ramsayi</i> Woma Python | P1 (SW subpop.) | - | Woodlands, heaths and shrublands, often with spinifex. Shelters mainly in abandoned monitor and mammal burrows and in soil cracks (Wilson and Swan, 2021). | One DBCA record 10.3 km northwest of the Survey Area in 2020 (DBCA, 2023c). | Medium |
| | | | | | | One record <15 km from the Survey Area, limited suitable habitat within the Survey Area. |
| Scincidae | <i>Ctenotus delli</i> Dell's Ctenotus | P4 | - | Restricted to the Darling range; Jarrah and Marri woodlands over shrubs on lateritic, sandy, and clay soils (Wilson and Swan, 2021). | Three DBCA records within 15 km of the Survey Area, including 11.6 km east of the Survey Area in 1981 (DBCA, 2023c). | High |
| | | | | | | Recorded <15 km from the Survey Area, and suitable habitat within the Survey Area. |
| Scincidae | <i>Ctenotus ora</i> Coastal Plain Skink | P3 | - | Open eucalypt woodland over <i>Banksia</i> and low vegetation on sandy coastal plain and coastal dunes south of Perth, from Pinjarra south to Yallingup Brook (Wilson and Swan, 2021). | One DBCA record 15 km west of the Survey Area in 2014 (DBCA, 2023c). | Low |
| | | | | | | No suitable habitat. |



5.0 Discussion

5.1 Fauna Habitat

The two fauna habitats, Jarrah-Marri Forest and Drainage Line recorded within the Survey Area, are typical of the Jarrah Forest bioregion. These habitats are consistent with the habitats previously recorded in studies in the region (**Appendix B**) and extend outside the Survey Area to form larger ecosystems which are widespread within the surrounding region. Overall connectivity at a regional level is relatively intact, as the Survey Area occurs within a landscape that retains large areas of natural vegetation. At a local level, habitats within and surround the immediate vicinity of the Survey Area are fragmented due to mining activities.

The Drainage Line habitat provides the most value to significant fauna and the overall fauna assemblage within the Survey Area. It is core habitat for all three targeted fauna taxa and often has surface water present which is a vital resource for the local fauna assemblage. A wide variety of fauna taxa will use this habitat as a dispersal corridor due the linear nature of the habitat and the presence of dense understory vegetation. Drainage Lines require relatively uninterrupted connectivity to allow fauna movement and dispersal, therefore, the Drainage Line habitat within the Survey Area is important to maintain this ecosystem function.

The Jarrah-Marri Forest habitat is relatively widespread and abundant at a regional scale. It is core habitat for the Chuditch and Woylie, and supporting habitat for the Quokka which tends to occupy dense understorey vegetation associated with wet areas such as Drainage Line habitat.

Both habitats contain tree hollows and hollow logs which are important microhabitats for hollow-dependent species. Hollow logs are often used by Chuditch for denning. The canopy of both habitats comprises valuable foraging and roosting habitat for birds such as black cockatoos, and the multitude of microhabitats (leaf litter, hummocks, woody debris, logs, and stones) provide refuge and feeding opportunities for a wide array of fauna species, particularly herpetofauna and small mammals.

5.2 Significant Fauna

5.2.1 Recorded within the Survey Area

Baudin's Cockatoo (*Zanda baudinii*) – EN (BC Act; EPBC Act)

The Baudin's Cockatoo is distributed through the south-west WA from the northern Darling Range and adjacent far east of the Swan Coastal Plain (south of the Swan River), south to Bunbury and across to Albany (Johnstone and Kirkby, 2011). The taxon rarely occurs near coastal communities north of Mandurah, and rarely occurs north of the Swan River (Johnstone and Storr, 1998; Johnstone and Kirkby, 2008). The species forages primarily in eucalypt forest, where it feeds on Marri seeds, flowers, nectar, and buds. They also feed on a wide range of seeds of *Eucalyptus* spp., *Banksia* spp., *Hakea* spp., and *Pinus* spp., as well as fruiting apples and pears and beetle larvae from under the bark of trees (Johnstone and Storr, 1998; Johnstone and Kirkby, 2008). Baudin's Cockatoo forages at all levels of the forest from the canopy to the ground, often feeding in the understorey on proteaceous trees and shrubs (especially *Banksia*) and in orchard trees within the canopy and on fallen fruit.

Secondary evidence of Baudin's Cockatoo was recorded during the survey, however direct sightings did not occur. The species can be cryptic due to its similarity in appearance to Carnaby's Cockatoos, and it can often be misidentified as such (Birdlife Australia, 2016). Within the Survey Area, the Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for the Baudin's Cockatoo because of their value for breeding, roosting, and foraging.



The taxon is likely to use the Survey Area for nesting and raising chicks in the breeding season and may use the habitats during the wintering season as core roosting and foraging habitat.

Forest Red-tailed Black Cockatoo (*Calyptorhynchus banksii naso*) - VU (BC Act; EPBC Act)

The Forest Red-tailed Black Cockatoo is distributed through the south-west of WA from Gingin through the Darling Ranges to the south-west from Bunbury to Albany. The Forest Red-tailed Black Cockatoo inhabits dense Jarrah, Karri (*E. diversicolor*), and Marri forests and feeds primarily on the fruit of Marri and Jarrah trees (Johnstone and Kirkby, 1999).

The Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for the Forest Red-tailed Black Cockatoo because of their value for breeding, roosting, and foraging. The taxon is likely to use the Survey Area for nesting and raising chicks during the breeding season and may use the habitats during the wintering season as core roosting and foraging habitat.

Chuditch (*Dasyurus geoffroii*) – VU (BC Act; EPBC Act)

The Chuditch inhabits areas dominated by sclerophyll forest or drier woodland, heath, and mallee shrubland (Van Dyck and Strahan, 2008). The species is generally highly mobile and uses bush remnants as corridors (Woinarski, Burbidge and Harrison, 2012). The Chuditch is a largely nocturnal animal, feeding on a carnivorous diet of mammals, birds, lizards, and frogs. They have been recorded foraging during the day such as during the breeding season or when cold and wet weather restricts their nocturnal movements (Van Dyck and Strahan, 2008). Most diurnal denning sites in sclerophyll forest consist of hollow logs or earth burrows, although Quenda nests and hollow tree bases may be used (Van Dyck and Strahan, 2008). The Chuditch was abundant prior to European settlement, and it is now largely restricted to the south-west of Western Australia, with small numbers in the Midwest, Wheatbelt and South Coast Regions (DBCA, 2017a).

The Chuditch was recorded via secondary evidence in the form of scat. The Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for the Chuditch because of their value for denning and foraging. The Drainage Line habitat is of particular importance as it provides water sources and dispersal corridors. Population numbers may be low and dispersed through the widespread Jarrah-Marri Forest making detection difficult.

Rakali (*Hydromys chrysogaster*) - P4 (DBCA)

The Rakali (also known as the Water Rat) is a native rodent species that lives in burrows on the banks of rivers, lakes, and estuaries and is an opportunistic predator feeding on aquatic insects, fish, crustaceans, mussels, snails, frogs, birds' eggs, and water birds (Atkinson, Lund, and Morris, 2008). The species can be found in permanent water systems in Australia, New Guinea, and offshore islands, with a home range of approximately 19 ha in coastal wetlands and streams of south-western Australia, individuals will establish sedentary territories where resident male and female ranges will overlap (Van Dyck and Strahan, 2008).

The Rakali was detected by camera trap within the Drainage Line habitat within the Survey Area. Drainage Line habitat constitutes core habitat for this species and the habitat within the survey area is important for connectivity.

Western Brush Wallaby (*Notamacropus irma*) – P4 (DBCA)

The Western Brush Wallaby occurs in the south-west of Western Australia and closely resembles a larger kangaroo (Van Dyck and Strahan, 2008). Its preferred habitat consists of open forest or woodland, particularly favouring open and seasonally wet flats with low grasses and open scrubby thickets (Van Dyck and Strahan, 2008). The Western Brush Wallaby was a common species during the early days of settlement, however, ongoing



clearing and fragmentation of bushland in the Wheatbelt as well as the dramatic increase in fox numbers within the south-west of Western Australia has led to this species' decline (Van Dyck and Strahan, 2008).

The Western Brush Wallaby was recorded during the survey by both camera trap and direct sighting. Within the Survey Area, the Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for the taxon because of their value for foraging and dispersal. The Drainage Line habitat is of particular importance as the taxon may rely on the seasonal wet flats within this habitat for the grasses and scrubby thickets for foraging and shelter.

5.2.2 High likelihood of occurrence within the Survey Area

Woylie (*Bettongia penicillata ogilbyi*) – CR (BC Act); EN (EPBC Act)

The Woylie is restricted to three small wheatbelt reserves in Western Australia: Dryandra Woodland, Tutanning Nature Reserve, and the Kingston/Perup Forest, all characterised by the presence thickets of the plant genus *Gastrolobium* (Van Dyck and Strahan, 2008). Fox predation is considered the driving factor in the Woylie's distribution contraction, with habitat destruction and disease also impacting populations to be rare and scattered (Van Dyck and Strahan, 2008). The Woylie primarily eats fungus and supplements its diet with seed kernels of the Desert Quandong (*Santalum acuminatum*), various tubers, bulbs, and invertebrates (Van Dyck and Strahan, 2008).

Within the Survey Area, the Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for the Woylie because of their value for shelter and foraging. The Drainage Line habitat is of particular significance to the taxon as this habitat contains valuable water sources and dense thickets which provide refuge from predators.

Carnaby's Cockatoo (*Zanda latirostris*) – EN (BC Act; EPBC Act)

Carnaby's Cockatoos nest in the hollows of a wide range of *Eucalyptus* trees, with a preference for smooth barked trees such as Salmon Gum (*E. salmonophloia*) and Wandoo (*E. wandoo*) but also rough barked *Eucalyptus* and *Corymbia* trees such as Red Morrell (*E. longicornis*), York Gum (*E. loxophleba*), Marri (*Corymbia calophylla*) and Tuart (*E. gomphocephala*) (Johnstone and Storr, 1998). Carnaby's Cockatoos feed on seeds, nuts, and flowers of a variety of native and exotic plants, including *Banksia* spp., Pine trees (*Pinus* sp.), Marri, Jarrah (*E. marginata*), *Grevillea* spp., *Allocasuarina* spp., and *Hakea* spp. (Shah, 2006).

The Jarrah-Marri Forest within the Survey Area provides core foraging, roosting and nesting habitat which may be used by this taxon. Recent previous studies have recorded foraging and unconfirmed breeding of this taxon in similar Jarrah-Marri Forest north of the Survey Area.

Quokka (*Setonix brachyurus*) – VU (BC Act; EPBC Act)

The Quokka is a small macropod found on two offshore islands (Rottnest and Bald Island) and several mainland sites in south-west Western Australia (Van Dyck and Strahan, 2008; DBCA, 2017a). The distribution of the Quokka is severely fragmented and there is little to no migration between populations. Mainland populations of the Quokka favour dense riparian vegetation; however, they also make use of heath and shrubland, swamp Peppermint, and swampy shrublands as well as Karri Forest regrowth (Van Dyck and Strahan, 2008; DBCA, 2017a). The populations of Quokkas on the mainland are largely restricted to areas of high rainfall of 1,000 mm or more (Van Dyck and Strahan, 2008).

The Drainage Line habitat constitutes core habitat for the Quokka due to the dense understorey vegetation. The Jarrah-Marri habitat constitutes supporting habitat for the taxon. A recent survey has recorded this taxon within Drainage Line habitat north of the Survey Area (GHD, 2021a). The Drainage Line habitat is limited in extent within the Survey Area



and extends outside the Survey Area, therefore it is possible that the taxon is present within the area but was not detected at camera trap locations during the survey.

Wambenger Brush-tailed Phascogale (*Phascogale tapoatafa wambenger*) – CD (BC Act)

The Wambenger Brush-tailed Phascogale is a rat-sized, arboreal, carnivorous marsupial, characterised by a tuft of black hair on its tail (Van Dyck and Strahan, 2008). Typically, the species has been observed in dry sclerophyll forests and open woodlands containing hollow-bearing trees with a sparse ground cover. The distribution of the Wambenger Brush-tailed Phascogale is widespread but fragmented, stemming from habitat destruction and predation by cats and foxes (Van Dyck and Strahan, 2008). In the south-west of Australia, the species is thought to have declined within the last 10 years given the lower number of records despite an increase in survey effort (Van Dyck and Strahan, 2008).

Wambenger Brush-tailed Phascogale was not recorded during the survey. The taxon has been recorded within 100 m of the Survey Area (DBCA, 2023c) and recent surveys in the region detected the species via trapping and camera trapping in nearby Jarrah-Marri Forest (GHD, 2021a). The Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for this species because of their value for shelter, foraging, and dispersal.

Quenda (*Isoodon fusciventer*) – P4 (DBCA)

Quenda are nocturnal and omnivorous, feeding on insects, spiders, worms, and plant roots and are typically found in dense vegetation, including wetland fringes, forest, woodland, shrub, and heath communities (Van Dyck and Strahan, 2008; DBCA, 2012). They are found only in the southwestern corner of Western Australia, where these regions generally display a combination of sandy soils with dense heathy vegetation in the lower stratum (Van Dyck and Strahan, 2008; DBCA, 2012).

Within the Survey Area, the Jarrah-Marri Forest and Drainage Line habitats constitute core habitat for the taxon because of their value for shelter, foraging, and dispersal. The Drainage Line habitat is of particular importance as provides water sources and dense understorey for shelter.

Western False Pipistrelle (*Falsistrellus mackenziei*) – P4 (DBCA)

The Western False Pipistrelle is endemic to the south-western region of Western Australia, making use of high rainfall forests dominated by Jarrah, Karri, Marri, and Tuart. The species is a specialist of tall mature forest, foraging within the canopy and roosting in tree hollows (Van Dyck and Strahan, 2008).

Western False Pipistrelle was not targeted during the surveys and so presence was not detected; however, this taxon is widely documented within the area and the Survey Area contains core Jarrah-Marri Forest and Drainage Line habitat which provides high value foraging habitat and roost sites which are associated with hollows in *Eucalyptus* (Baker and Gynther, 2023).

Dell's Ctenotus (*Ctenotus delli*) – P4 (DBCA)

Dell's Ctenotus is endemic to the north Darling Range, south-western Australia, inhabiting dry sclerophyll forest on stony hills and ranges (Cogger, 2014).

This taxon may use the Jarrah-Marri Forest with lateritic clay substrate within the Survey Area which provides suitable foraging and breeding habitats.



5.2.3 Medium likelihood of occurrence within the Survey Area

Western Ringtail Possum (*Pseudocheirus occidentalis*) – CR (BC Act; EPBC Act)

The Western Ringtail Possum is found in the South-West, particularly around Busselton, largely in areas of Peppermint (*Agonis flexuosa*) or Tuart (*Eucalyptus gomphocephala*) forest, generally with a Peppermint midstory (Van Dyck and Strahan, 2008). The species is nocturnal, and typically feeds on the leaves of Peppermint or Eucalypt trees. In coastal Peppermint habitat, Western Ringtail Possums constructs dreys anywhere from the upper canopy to the understory, which are used as rest sites during the day. Coastal populations are under immediate threat from habitat loss and fragmentation and recent research from Jarrah Forest sites indicates that increasing fire intensity at the local scale and fox predation are also impacting the species (Van Dyck and Strahan, 2008).

The Western Ringtail Possum has been recorded within the Desktop Study Area and may occur within the Survey Area. Records range between 1997 and 2020 due to multiple translocations occurring at Lane Poole Reserve (17 km north of the Survey Area), Yalgorup National Reserve (37 km northwest of the Survey Area), and Leschenault Peninsula (41 km southwest of the Survey Area) between 1991 and 2004. Due to predation, disease, habitat destruction by fires and limited post-translocation monitoring data, translocated populations were unstable with numbers considered to be low and undetectable. The Survey Area has the potential to support this taxon as Jarrah-Marri Forest constitutes core habitat for inland populations where Peppermint and Tuart is absent.

Australasian Bittern (*Botaurus poiciloptilus*) – EN (BC Act)

The Australasian Bittern is a secretive bird with a distinct booming call that nests and forages within dense wetland vegetation (Johnstone and Storr, 1998). In Western Australia, the Australasian Bittern was formerly widespread throughout the southwest, however this range has declined. It is now considered that the Australasian Bittern only occurs in select habitats, including the western coastal plain between Lancelin and Busselton, the southern coastal region from Augusta to the east of Albany and inland to some wetlands in the Jarrah Forest belt with small, isolated populations in swamps from west of Esperance eastwards to near Cape Arid (Marchant and Higgins, 1990). The main cause of this decline is thought to be wetland drainage and degradation.

Although the Australasian Bittern typically occurs in swamps, the taxon was recorded 5.8 km north of the Survey Area in 2017 in Drainage Line habitat that appears to be similar to that which occurs within the Survey Area. Therefore, the Drainage Line habitat constitutes potential habitat for the taxon.

Malleefowl (*Leipoa ocellata*) – VU (BC Act; EPBC Act)

The Malleefowl is found in semi-arid to arid shrublands and low woodlands, especially those dominated by mallee or *Acacia*. A sandy substrate and abundance of leaf litter are required for breeding (DEE, 2018). Densities of the birds are generally greatest in areas of higher rainfall and on more fertile soils where habitats tend to be thicker and there is an abundance of food plants. Much of the best habitat for Malleefowl has already been cleared or has been modified by grazing via Sheep, Cattle, Rabbits, and Goats (Benshemesh, 2007). The species nests in large mounds of dirt and leaf litter, up to five metres wide and one metre tall (Menkhorst *et al.*, 2019).

Malleefowl was sighted 17.6 km from the Survey Area in 2014 (DBCWA, 2023c) Furthermore, individuals were translocated to the nearby Dryandra Woodland, 79 km east-northeast of the Survey Area, within the last five years (Traill, 2019). The Jarrah-Marri Forest and Drainage Line habitats have the potential to support this species, though no evidence of foraging or nesting was recorded during the survey. Malleefowl are often cryptic and nest in areas of dense shrubland, making detection of the species difficult without LiDAR mapping of potential nesting mounds.



Peregrine Falcon (*Falco peregrinus*) – OS (DBCA)

The Peregrine Falcon is an uncommon but wide-ranging bird across Australia (Barrett *et al.*, 2003). It occurs along rivers and ranges as well as wooded watercourses and lakes. It nests primarily on cliffs, granite outcrops and quarries, although is also known to occupy existing raptor and corvid stick nests (Menkhorst *et al.*, 2019). The diet of the Peregrine Falcon has been well studied and primarily includes flocking species such as parrots, pigeons and on the east coast, European Starlings (Olsen and Fuentes, 2008).

Peregrine Falcon was not observed during the survey; however, this species has been previously recorded within the region flying over similar Jarrah-Marri Forest (GHD, 2021a). In the absence of cliff ledges, Peregrine Falcons will use trees for nesting (Morcombe, 2003). The Jarrah-Marri Forest within the Survey Area and surrounds may provide suitable nesting and hunting habitat to support this species.

Woma Python (*Aspidites ramsayi*) – P1 (DBCA)

The Woma Python is a nocturnal, terrestrial snake that inhabits woodland, heaths, and shrublands, often with spinifex. It shelters in hollow logs, abandoned monitor and mammal burrows, and soil cracks (Wilson and Swan, 2021). It is usually restricted to arid and dry inland areas, however there is an isolated population in south-western Australia (Cogger, 2014).

Within the Survey Area, both the Jarrah-Marri Forest and Drainage Line habitats provide suitable resources for hunting and shelter.

Southern Death Adder (*Acanthophis antarcticus*) – P4 (DBCA)

The Southern Death Adder is found in Southwest and South Australia to Eastern and Central Queensland where, due to habitat destruction and altered fire regimes, populations are in decline (Cogger, 2014; Wilson and Swan, 2021). Habitats range from rainforest to shrublands and heath (Wilson and Swan, 2021).

The Survey Area falls within distribution of the Southern Death Adder and microhabitats within the Jarrah-Marri Forest and Drainage Line habitats have the potential to be used by the taxon for shelter and hunting.



6.0 Conclusion

Two fauna habitats were mapped within the Survey Area, Jarrah-Marri Forest, and Drainage Line habitats. The Drainage Line habitat provides the most value for the targeted taxa as it is core habitat for the Chuditch, Quokka, and Woylie. The Jarrah-Marri Forest also provides value for targeted taxa as core habitat for the Chuditch and Woylie and supporting habitat for the Quokka.

One targeted taxon was recorded during the survey; secondary evidence of Chuditch was recorded in the form of scat within the Jarrah-Marri Forest. The Quokka and Woylie were not recorded during the current survey however, they have previously been recorded within 200 m and 40 m of the Survey Area respectively.

In addition to the targeted fauna taxa, four significant taxa were recorded during the fauna survey:

- Forest Red-tailed Black Cockatoo – VU (EPBC; BC Act).
- Baudin's Cockatoo – EN (EPBC; BC Act).
- Western Brush Wallaby – P4 (DBCA).
- Rakali – P4 (DBCA).

A further seven significant fauna taxa were assessed as having a high likelihood and six significant fauna taxa were assessed as having a medium, and 15 significant fauna taxa were assessed as having have a low likelihood of occurring within the Survey Area.

Three introduced taxa were recorded during the survey: Pig (*Sus scrofa*), Cat (*Felis catus*) and Black Rat (*Rattus rattus*).



7.0 References

- Atkinson, C., Lund, M. and Morris, K. (2008) 'BiblioRakali: the Australian water rat, *Hydromys chrysogaster* Geoffroy, 1804 (Muridae: Hydromyinae), a subject-specific bibliography', *Conservation Science W. Aust.*, 7(1), pp. 65–71.
- Baker, A.M. and Gynther, I.C. (2023) *Strahan's Mammals of Australia*. 4th edn. Wahroonga, Australia: Reed New Holland Publishers.
- Barrett, G. *et al.* (2003) *The new atlas of Australian birds*. Melbourne, Australia: Royal Australasian Ornithologists Union.
- Beard, J.S. *et al.* (2013) 'The vegetation of Western Australia at the 1:3,000,000 scale. Explanatory memoir. Second edition', *Conservation Science W. Aust.*, 9(1), pp. 1–152.
- Benshemesh, J. (2007) *National recovery plan for malleefowl *Leipoa ocellata**. Adelaide, Australia.
- Birdlife Australia (2016) *Identifying Southwest Black-Cockatoos*. Perth, Australia.
- BoM (2007) *About Climate Statistics*. Available at: <http://www.bom.gov.au>.
- BoM (2024) *Climate Data Online*. Available at: <http://www.bom.gov.au>.
- Cogger, H.G. (2014) *Reptiles and amphibians of Australia*. 7th edn. Melbourne, Australia: CSIRO Publishing.
- Commonwealth of Australia (1999) *Environment Protection and Biodiversity Conservation Act 1999*. Australia. Available at: www.legislation.gov.au.
- DBCA (2012) *Fauna Profile - Quenda *Isoodon obesulus**. Available at: www.dpaw.wa.gov.au.
- DBCA (2017a) *Fauna Profile - Chuditch *Dasyurus geoffroii**. Available at: <http://www.dbca.wa.gov.au>.
- DBCA (2017b) *Fauna Profile - Quokka *Setonix brachyurus**. Available at: <https://www.dpaw.wa.gov.au/>.
- DBCA (2023a) *DBCA - Lands of Interest (DBCA-012)*. Available at: <https://catalogue.data.wa.gov.au>.
- DBCA (2023b) *DBCA - Legislated Lands and Waters (DBCA-011)*. Available at: <https://catalogue.data.wa.gov.au>.
- DBCA (2023c) *DBCA Threatened and Priority Fauna Database request (Custom Search)*.
- DBCA (2023d) *NatureMap Database Search*. Perth, Australia.
- DCCEEW (2023) *Protected Matters Search Tool*. Available at: <https://www.dcceew.gov.au/environment/epbc/protected-matters-search-tool>.
- DEE (2016) *Interim Biogeographic Regionalisation for Australia, Version 7*. Canberra, Australia. Available at: www.environment.gov.au.
- DEE (2018) *Leipoa ocellata - Malleefowl*. Available at: http://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon_id=934.
- DMIRS (2024) *Mining Tenements (DMIRS-003)*. Available at: <https://catalogue.data.wa.gov.au/>.
- DoE (2013) *Matters of National Environmental Significance Significant impact guidelines 1.1 Environment Protection and Biodiversity Conservation Act 1999*. Canberra, Australia. Available at: <https://www.environment.gov.au>.
- DPIRD (2022) *Soil Landscape Mapping - Best Available (DPIRD-027)*. Perth, Australia. Available at: <https://catalogue.data.wa.gov.au>.



- DSEWPaC (2011) *Survey guidelines for Australia's threatened mammals: Guidelines for detecting mammals listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999*. Canberra, Australia. Available at: <https://www.dcceew.gov.au>.
- DWER (2018) *Hydrography, Linear (Hierarchy) (DWER-031)*. Perth, Australia: Landgate. Available at: <https://catalogue.data.wa.gov.au>.
- DWER (2021) *Clearing Regulations - Environmentally Sensitive Areas (DWER-046)*. Available at: <https://catalogue.data.wa.gov.au/>.
- Van Dyck, S. and Strahan, R. (2008) *The mammals of Australia*. 3rd edn. Sydney, Australia: New Holland Publishers.
- Environmental Management and Research Consultants (2006) *Vertebrate Fauna Survey of Planned Mining Areas at Alcoa's Willowdale Larego Mining Region*.
- EPA (2016) *Environmental Factor Guideline - Terrestrial Fauna*. Perth. Available at: <https://www.epa.wa.gov.au>.
- EPA (2020) *Technical Guidance – Terrestrial vertebrate fauna surveys for environmental impact assessment*. Available at: <https://www.epa.wa.gov.au>.
- GHD (2012) *Report for Rail Reserves in the Shire of Serpentine Jarrahdale - Spring Flora and Vegetation Survey and Fauna and Habitat Assessment*.
- GHD (2021a) *Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine-Holyoake*.
- GHD (2021b) *Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine-Myara North*.
- Government of Western Australia (1986) *Environmental Protection Act 1986*. Available at: www.legislation.wa.gov.au.
- Government of Western Australia (2016) *Biodiversity Conservation Act 2016*. Available at: www.legislation.wa.gov.au.
- Harewood, G. (2017) *Fauna Assessment Pinjarra - Williams Road (SLK 14 – 67)*.
- Harewood, G. (2019) *Black Cockatoo Habitat Assessment - Pinjarra Racecourse*.
- Johnstone, R.E. and Kirkby, T. (1999) 'Food of the forest red-tailed black cockatoo *Calyptorhynchus banksii naso* in south-west Western Australia', *Western Australian Naturalist*, 22, pp. 167–177.
- Johnstone, R.E. and Kirkby, T. (2011) *Carnaby's Black Cockatoo (*Calyptorhynchus latirostris*), Baudin's Black Cockatoo (*Calyptorhynchus baudinii*) and the Forest Red-tailed Black Cockatoo (*Calyptorhynchus banksii naso*) on the Swan Coastal Plain (Lancelin–Dunsborough), Western Australia*. Perth, Australia.
- Johnstone, R.E. and Storr, G.M. (1998) *Handbook of Western Australian birds*. Edited by D. Louise. Perth, Australia: Western Australian Museum.
- Marchant, S. and Higgins, P.J. (1990) *Handbook of Australian, New Zealand and Antarctic Birds. Volume 1 - Ratites to Ducks*. Melbourne: Oxford University Press.
- Menkhorst, P. *et al.* (2019) *The Australian Bird Guide*. Revised Edition. Melbourne, Australia: CSIRO Publishing.
- Morcombe, M. (2003) *Field Guide to Australian Birds*. Archerfield: Steve Parish Publishing Pty Ltd.
- Olsen, J. and Fuentes, E. (2008) 'Dietary shifts based upon prey availability in Peregrine Falcons and Australian Hobbies breeding near Canberra, Australia', *Journal of Raptor Research*, 42, pp. 125–137.



Pizzey, G. and Knight, F. (2001) *Field Guide to Birds of Australia*. Pymble, Australia: HarperCollins Australia.

Shah, B. (2006) *Conservation of Carnaby's Black Cockatoo on the Swan Coastal Carnaby's plain, Western Australia*. Perth, Australia.

Shepherd, D.P., Beeston, G.R. and Hopkins, A.J. (2002) *Native vegetation in Western Australia: Extent, type and status. Resource Management Technical Report 249*. Perth, Australia. Available at: <https://library.dpird.wa.gov.au>.

Trill, M. (2019) 'Malleefowl land in Dryandra', *Narrogin Observer*. Available at: <https://www.narroginobserver.com.au>.

Way, S. (2013) *Chuditch Pre-mine Survey, Myara* .

Williams, K. and Mitchell, D. (2001) *Jarrah Forest 1 (JF1 – Northern Jarrah Forest subregion)*. Perth, Australia.

Wilson, S. and Swan, G. (2021) *A complete guide to reptiles of Australia*. 6th edn. New Holland Publishers.

Woinarski, J., Burbidge, A. and Harrison, P. (2012) *The action plan for Australian mammals 2012*. CSIRO Publishing.



8.0 Feedback

At SLR, we are committed to delivering professional quality service to our clients. We are constantly looking for ways to improve the quality of our deliverables and our service to our clients. Client feedback is a valuable tool in helping us prioritise services and resources according to our client needs.

To achieve this, your feedback on the team's performance, deliverables and service are valuable and SLR welcome all feedback via <https://www.slrconsulting.com/en/feedback>. We recognise the value of your time, and we will make a \$10 donation to our 2023 Charity Partner - Lifeline, for every completed form.





Appendix A Maps

Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd

SLR Project No.: 675.VX5890.00001

17 June 2024

410000

420000

6360000



6360000

6350000


6350000

6340000

6340000



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:100,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 18/03/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Area
 MAP 1

410000


415000

6355000

6355000

6350000

6350000



| | |
|-------------------------------|-------------|
| | Survey Area |
| | Road |
| Soil Landscape Mapping | |
| | 255DpDW |
| | 255DpHR |
| | 255DpMH |
| | 255DpYG |
| | 255DpYGh |
| | 255DpYGsw |



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:50,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 18/03/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Soil Landscape Mapping
 MAP 2

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.

410000


415000

6355000

6355000

6350000

6350000



| | |
|--|--------------------|
| | Survey Area |
| | Road |
| | Minor River |
| | Mainstream |
| | Major Tributary |
| | Significant Stream |



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:50,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 18/03/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Hydrography
 MAP 3

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.

410000


415000

6355000

6355000

6350000

6350000




Legend

- Survey Area
- Road
- Pre European Vegetation
- West Darling 3



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:50,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 18/03/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Pre-European Vegetation
 MAP 4


COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:75,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 16/04/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Conservation and Environmentally
 Sensitive Areas
 MAP 5



| I.D. | PREVIOUS SURVEY |
|------|--|
| 1 | Report for Rail Reserves in the Shire of Serpentine Jarrahdale - Spring Flora and Vegetation Survey and Fauna and Habitat Assessment (GHD, 2012) |
| 2 | Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Myara North (GHD, 2021b) |
| 3 | Chuditch Pre-mine Survey, Myara (Way, 2013) |
| 4 | Vertebrate Fauna Survey of Planned Mining Areas at Alcoa's Willowdale Larego Mining Region (Environmental Management and Research Consultants, 2006) |
| 5 | Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Holyoake (GHD, 2021a) |
| 6 | Fauna Assessment Pinjarra - Williams Road (SLK 14 fC6 67) (Harewood, 2017) |



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:500,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 18/03/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region


Previous Surveys in the Locality
 MAP 6

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.

410000

415000

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



635000

635000


635000

635000

Survey Area
 GPS Tracks
 Camera Trap



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.







Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:50,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-1

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



-  Survey Area
-  Survey Track
-  Camera Trap




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-2


COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



Survey Area
 Survey Track
 Camera Trap



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-3


COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



| | |
|--|--------------|
| | Survey Area |
| | Survey Track |
| | Camera Trap |



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-4

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.








Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-5

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.

-  Survey Area
-  Survey Track
-  Camera Trap




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-6

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT. THE COMMISSIONER, SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



| | |
|---|--------------|
|  | Survey Area |
|  | Survey Track |
|  | Camera Trap |




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-7

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



-  Survey Area
-  Survey Track
-  Camera Trap







DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-8

-  Survey Area
-  Survey Track
-  Camera Trap



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD




Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region


Survey Effort
 MAP 7-9

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



-  Survey Area
-  Survey Track
-  Camera Trap



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-10



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.


COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



Survey Area
 Survey Track
 Camera Trap



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.







Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
 MAP 7-11


COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



-  Survey Area
-  Survey Track
-  Camera Trap



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 14/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Survey Effort
MAP 7-12


400000



| | |
|--|---|
| Survey Area | Specially Protected - Conservation Dependent |
| Road | Specially Protected - Migratory |
| Significant Fauna Database Search Results | Specially Protected - Other Specially Protected |
| Threatened - Critically Endangered | Priority 1 |
| Threatened - Endangered | Priority 3 |
| Vulnerable | Priority 4 |



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.




Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:250,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 18/03/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region

Significant Fauna Database Search Results
 MAP 8

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.




COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR WHICH IT WAS COMMISSIONED. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.

6350000

6350000

Legend

- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- ★ Western Brush Wallaby (P4)
- + Rakali (P4)




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:50,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
MAP 9-1

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



Legend

- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- ★ Western Brush Wallaby (P4)
- + Rakali (P4)











DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.


Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-2

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



-  Survey Area
- Fauna Habitat
-  Jarrah Marri Forest
-  Drainage Line
- Significant Fauna Record
-  Baudin's Cockatoo (EN)
-  Forest Red-tailed Black Cockatoo (VU)
-  Western Brush Wallaby (P4)
-  Rakali (P4)



Service Layer Credits: Landgate / SLIP

Coordinate System: GDA 1994 MGA Zone 50 @ A4

Scale : 1:10,000 @ A4

Project Number : 675.VX5890.00001

Date Drawn : 7/06/2024

Drawn By : Environmaps


Reviewed By : PD




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-3

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- Western Brush Wallaby (P4)
- Rakali (P4)



Service Layer Credits: Landgate / SLIP


Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-4

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- Western Brush Wallaby (P4)
- Rakali (P4)




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.


Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-5

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- Western Brush Wallaby (P4)
- Rakali (P4)



Service Layer Credits: Landgate / SLIP


Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.


Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-6

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



Survey Area

Fauna Habitat

Jarrah Marri Forest

Drainage Line

Significant Fauna Record

Baudin's Cockatoo (EN)


Forest Red-tailed Black Cockatoo (VU)

Western Brush Wallaby (P4)

Rakali (P4)



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4

Scale : 1:10,000 @ A4

Project Number : 675.VX5890.00001

Date Drawn : 7/06/2024

Drawn By : Environmaps

Reviewed By : PD

Alcoa


Targeted Chuditch, Quokka, and Woylie Surveys Willowdale Mine

Larego Region

Fauna Habitat and Significant Fauna Records


MAP 9-8

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



Legend

- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- ★ Western Brush Wallaby (P4)
- + Rakali (P4)




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.


Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-9

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



| | |
|---------------------------------|---------------------------------------|
| | Survey Area |
| Fauna Habitat | |
| | Jarrah Marri Forest |
| | Drainage Line |
| Significant Fauna Record | |
| | Baudin's Cockatoo (EN) |
| | Forest Red-tailed Black Cockatoo (VU) |
| | Western Brush Wallaby (P4) |
| | Rakali (P4) |




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.

Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-10

COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.




Legend

- Survey Area
- Fauna Habitat**
- Jarrah Marri Forest
- Drainage Line
- Significant Fauna Record**
- Baudin's Cockatoo (EN)
- Forest Red-tailed Black Cockatoo (VU)
- ★ Western Brush Wallaby (P4)
- + Rakali (P4)




DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.










Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD

Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
MAP 9-11


COPYRIGHT: THIS DOCUMENT IS AND SHALL REMAIN THE PROPERTY OF SLR CONSULTING. THIS DOCUMENT MAY ONLY BE USED FOR THE PURPOSE FOR WHICH IT WAS COMMISSIONED AND IN ACCORDANCE WITH THE TERMS OF ENGAGEMENT FOR THE COMMISSION. SLR CONSULTING DOES NOT HOLD ANY RESPONSIBILITY FOR THE MISUSE OF THIS DOCUMENT.



-  Survey Area
- Fauna Habitat**
-  Jarrah Marri Forest
-  Drainage Line
- Significant Fauna Record**
-  Baudin's Cockatoo (EN)
-  Forest Red-tailed Black Cockatoo (VU)
-  Western Brush Wallaby (P4)
-  Rakali (P4)



DISCLAIMER: All information within this document may be based on external sources. SLR Consulting Pty Ltd makes no warranty regarding data's accuracy or reliability for any purpose.



Coordinate System: GDA 1994 MGA Zone 50 @ A4
 Scale : 1:10,000 @ A4
 Project Number : 675.VX5890.00001
 Date Drawn : 7/06/2024
 Drawn By : Environmaps
 Reviewed By : PD



Alcoa
 Targeted Chuditch, Quokka, and
 Woylie Surveys Willowdale Mine
 Larego Region
 Fauna Habitat and
 Significant Fauna Records
 MAP 9-12



Appendix B Literature Review Summary

Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd

SLR Project No.: 675.VX5890.00001

17 June 2024

| ID | Report | Survey location | Survey timing | Survey type | Significant fauna recorded during the survey | Fauna habitats recorded during the survey |
|-------|---|--------------------------------|---------------|---|--|---|
| Lit A | <i>Fauna Assessment Pinjarra - Williams Road (SLK 14 – 67)</i> (Harewood, 2017) | 14 km north of the Survey Area | August 2011 | Basic Fauna Assessment and Black Cockatoo Habitat Assessment | <ul style="list-style-type: none"> • Baudin's Cockatoo (<i>Zanda baudinii</i>) – Listed as Endangered under the BC Act and EPBC Act • Carnaby's Cockatoo (<i>Zanda laltrostris</i>) - Listed as Endangered under the BC Act and EPBC Act • Forest Red-tailed Cockatoo (<i>Calyptorhynchus banksii naso</i>) – Listed at Vulnerable under the BC Act and EPBC Act • Western Brush Wallaby (<i>Notamacropus irma</i>) - Listed as Priority 4 by the DBCA | <p>Four fauna habitats were identified:</p> <ul style="list-style-type: none"> • Remnant native vegetation • Plantations • Existing cleared areas • River and ephemeral streams |
| Lit B | <i>Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Holyoake</i> (GHD, 2021a) | 18 km north of the Survey Area | August 2021 | Targeted Forest Red-tailed Black Cockatoo, Carnaby's Cockatoo, Baudin's Cockatoo, Chuditch, Quokka, Carters's Freshwater Mussel, and other Priority species | <ul style="list-style-type: none"> • Baudin's Cockatoo (<i>Zanda baudinii</i>) – Listed as Endangered under the BC Act and EPBC Act • Carnaby's Cockatoo (<i>Zanda laltrostris</i>) - Listed as Endangered under the BC Act and EPBC Act • Forest Red-tailed Cockatoo (<i>Calyptorhynchus banksii naso</i>) – Listed at Vulnerable under the BC Act and EPBC Act | <p>Seven fauna habitats were identified:</p> <ul style="list-style-type: none"> • Bullich forest • Granite outcrop • Blackbutt forest • Flooded Gum woodland • Jarrah-Marri forest • Mine rehabilitation • Pine plantation |

| ID | Report | Survey location | Survey timing | Survey type | Significant fauna recorded during the survey | Fauna habitats recorded during the survey |
|-------|---|--------------------|---------------|--|---|--|
| | | | | | <ul style="list-style-type: none"> • Quokka (<i>Setonix brachyurus</i>) – Listed as Vulnerable under the BC Act and EPBC Act • Chuditch (<i>Dasyurus geoffroii fortis</i>) - Listed at Vulnerable under the BC Act and EPBC Act • Peregrine Falcon (<i>Falco peregrinus</i>) – Listed as Special Protection under the BC Act • Brush-tailed Phascogale (<i>Phascogale tapoatafa wambenger</i>) – Listed as Conservation Dependent By DBCA • Western False Pipistrelle (<i>Falsistrellus mackenziei</i>) - Listed as Priority 3 by the DBCA • Quenda (<i>Isodon fusciventer</i>) - Listed as Priority 4 by the DBCA • Western Brush Wallaby (<i>Notamacropus irma</i>) - Listed as Priority 4 by the DBCA | |
| Lit C | <i>Black Cockatoo Habitat Assessment – Pinjarra Racecourse</i> (Harewood, 2019) | 30 km north of the | 2019 | Targeted Black Cockatoo Habitat Assessment | <ul style="list-style-type: none"> • Forest Red-tailed Cockatoo (<i>Calyptorhynchus banksii naso</i>) – Listed at | No fauna habitats were identified. However, 34 habitat trees consisting of Flooded Gum, Tuart, |

| ID | Report | Survey location | Survey timing | Survey type | Significant fauna recorded during the survey | Fauna habitats recorded during the survey |
|-------|---|--------------------------------|--------------------|--|---|--|
| | | Survey Area | | | Vulnerable under the BC Act and EPBC Act | Marri and introduces eucalypts were identified. |
| Lit D | <i>Chuditch Pre-mine Survey, Myara</i> (Way, 2013) | 43 km north of the Survey Area | March – April 2013 | Targeted Chuditch Survey | <ul style="list-style-type: none"> • Chuditch (<i>Dasyurus geoffroii fortis</i>) - Listed at Vulnerable under the BC Act and EPBC Act | Two fauna habitats were identified: <ul style="list-style-type: none"> • Jarrah forest • Drainage line |
| Lit E | <i>Vertebrate Fauna Survey of Planned Mining Areas at Alcoa's Willowdale Larego Mining Region</i> (Environmental Management and Research Consultants, 2006) | 43 km north of the Survey Area | 1996-2006 | Long Term Fauna Monitoring Program | <ul style="list-style-type: none"> • Baudin's Cockatoo (<i>Zanda baudinii</i>) – Listed as Endangered under the BC Act and EPBC Act • Noisy Scrubbird (<i>Atrichornis clamosus</i>) – Listed as Endangered under the BC Act and EPBC Act • Quokka (<i>Setonix brachyurus</i>) – Listed as Vulnerable under the BC Act and EPBC Act | No fauna habitats were recorded. |
| Lit F | <i>Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Myara North</i> (GHD, 2021b) | 54 km north of Survey Area | August 2021 | Targeted Forest Red-tailed Black Cockatoo, Carnaby's Cockatoo, Baudin's Cockatoo, Chuditch, Quokka, and other Priority species | <ul style="list-style-type: none"> • Baudin's Cockatoo (<i>Zanda baudinii</i>) – Listed as Endangered under the BC Act and EPBC Act • Carnaby's Cockatoo (<i>Zanda laltrostris</i>) - Listed as Endangered under the BC Act and EPBC Act • Forest Red-tailed Cockatoo (<i>Calyptorhynchus banksii naso</i>) – Listed at | Eight fauna habitats were identified: <ul style="list-style-type: none"> • Jarrah-Marri forest • Bullich forest • Granite outcrop • Blackbutt forest • Flooded Gum woodland • Melaleuca dampland • Mine rehabilitation • Pine plantation |

| ID | Report | Survey location | Survey timing | Survey type | Significant fauna recorded during the survey | Fauna habitats recorded during the survey |
|----|--------|-----------------|---------------|-------------|--|---|
| | | | | | <p>Vulnerable under the BC Act and EPBC Act</p> <ul style="list-style-type: none"> • Quokka (<i>Setonix brachyurus</i>) – Listed as Vulnerable under the BC Act and EPBC Act • Chuditch (<i>Dasyurus geoffroii fortis</i>) - Listed at Vulnerable under the BC Act and EPBC Act • Peregrine Falcon (<i>Falco peregrinus</i>) – Listed as Special Protection under the BC Act • Brush-tailed Phascogale (<i>Phascogale tapoatafa wambenger</i>) – Listed as Conservation Dependent By DBCA • Masked Owl Southwest (<i>Tyto novaehollandiae</i>) – Listed as Priority 3 by the DBCA • Southern Death Adder (<i>Acanthophis antarcticus</i>) – Listed as Priority 3 by the DBCA • Rakali (<i>Hydromys chrysogaster</i>) – Listed as Priority 4 by the DBCA | |

| ID | Report | Survey location | Survey timing | Survey type | Significant fauna recorded during the survey | Fauna habitats recorded during the survey |
|-------|--|--------------------------------|----------------|--------------------------------------|---|--|
| | | | | | <ul style="list-style-type: none"> • Quenda (<i>Isoodon fusciventer</i>) - Listed as Priority 4 by the DBCA • Western Brush Wallaby (<i>Notamacropus irma</i>) - Listed as Priority 4 by the DBCA • Western False Pipistrelle (<i>Falsistrellus mackenziei</i>) - Listed as Priority 3 by the DBCA | |
| Lit G | <p><i>Report for Rail Reserves in the Shire of Serpentine Jarrahdale - Spring Flora and Vegetation Survey and Fauna and Habitat Assessment</i> (GHD, 2012)</p> | 60 km north of the Survey Area | November, 2011 | Level 1 Fauna and Habitat Assessment | <ul style="list-style-type: none"> • Forest Red-tailed Cockatoo (<i>Calyptorhynchus banksii naso</i>) – Listed at Vulnerable under the BC Act and EPBC Act • Baudin’s Cockatoo (<i>Zanda baudinii</i>) – Listed as Endangered under the BC Act and EPBC Act • Western Brush Wallaby (<i>Notamacropus irma</i>) - Listed as Priority 4 by the DBCA • Quenda (<i>Isoodon fusciventer</i>) - Listed as Priority 4 by the DBCA • Letter-winged kite (<i>Elanus scriptus</i>) – Listed as Priority 4 the DBCA | <p>Five fauna habitats were identified:</p> <ul style="list-style-type: none"> • Melaleuca wetlands • Swampland • Jarrah/Marri forest • <i>Banksia/Allocasuarina</i> woodlands • Dense shrublands on clay flats |



Appendix C Licences and Permits

Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd

SLR Project No.: 675.VX5890.00001

17 June 2024



Wildlife Conservation Act 1950
REGULATION 17

Fauna taking (scientific or other purposes) licence (Fauna taking (scientific or other purposes) licence)

The undermentioned person may take fauna for research or other scientific purposes and where authorised, keep it in captivity, subject to the following and attached conditions, which may be added to, suspended or otherwise varied as considered fit.

Director General

Conditions

- 1 The licensee must comply with the provisions of the Wildlife Conservation Act 1950, Wildlife Conservation Regulations 1970 and any Notices in force under this legislation.
- 2 The licensee shall take fauna only in the manner stated on the endorsed Regulation 17 licence application form and endorsed related correspondence.
- 3 Unless specifically authorised in the conditions of this Licence or otherwise in writing by the Director General, species of fauna declared as likely to become extinct, rare or otherwise in need of special protection shall not be taken.
- 4 Any by-catch of fauna, which is declared to be rare, likely to become extinct, or otherwise in need of special protection shall be released immediately at the point of capture. Where such fauna taken under this licence is injured or deceased, the licensee shall contact the Department's Wildlife Licensing Section for advice on disposal. Records must be kept of any such fauna so captured and details are to be included in the report required under further condition below.
- 5 Any interaction involving Gazetted Threatened Fauna that may be harmful to the fauna and/or invasive may require approval from the Commonwealth Department of the Environment ph 02 6274 1111. Interaction with such species is controlled by the Commonwealth Environment Protection and Biodiversity Conservation Act 1999 and Environment Protection and Biodiversity Conservation Regulations 2000 as well as the Wildlife Conservation Act 1950 and Wildlife Conservation Regulations 1970.
- 6 No fauna shall be taken in areas where it would impinge on pre-existing scientific research programs.
- 7 Except in the case of approved lethal traps, the licensee shall ensure that measures are taken in the capture and handling of fauna to prevent injury or mortality resulting from that capture or handling. Where traps or other mechanical means or devices are used to capture fauna these shall be deployed so as to prevent exposure of trapped animals to ants and debilitating weather conditions and inspected at regular intervals throughout each day of their use. At the conclusion of research all markers used, and signs and structures erected by the licensee shall be removed and the environment returned to its original condition.
- 8 Not more than ten specimens of any one protected species of fauna shall be taken and removed from any location less than 20km apart. Where exceptional circumstances make it necessary to take a larger number of specimens from a particular location in order to obtain adequate statistical data, the collector must proceed with circumspection and justify their actions to the Director General in advance.
- 9 The licensee shall not release any fauna or their progeny in any area where it does not naturally occur, nor hand such fauna over to any other person or authority unless approved by the Director General, nor dispose of the remains of such fauna in any manner likely to confuse the natural or present day distribution of the species.
- 10 Bioprospecting involving the removal of sample aquatic and terrestrial organisms for chemical extraction and bioactivity screening shall not be conducted without specific written approval by the Director General.
- 11 No fauna is to be taken from any CALM land, as defined in the Conservation and Land Management Regulations 2002, without prior written approval of the Director General. No fauna is to be taken from any public land without the prior written approval of the Government Authority managing that land.
- 12 The licensee must not enter upon any private property or pastoral lease for the purposes of this licence, nor take any fauna from any private land or pastoral lease without the prior consent in writing of the owner or occupier. Similarly, in the case of Aboriginal lands, the licensee must not enter upon or take fauna from such lands without the written approval of the Department of Aboriginal Affairs and/or the relevant native title holders or applicants.
- 13 Copies of this licence and any written approval or consent required by conditions of this licence must be carried by the licensee and any person/s authorised under the licence at all times when conducting activities relevant to the licence

DEPARTMENT OF PARKS AND WILDLIFE



Department of
Parks and Wildlife



Enquiries: 17 DICK PERRY AVE, KENSINGTON, WESTERN AUSTRALIA
Telephone: 08 9219 9000
Facsimile: 08 9219 8242
Web Site: <https://wildlifelicensing.dpaw.wa.gov.au>
Correspondance: **Locked Bag 30**
Bentley Delivery Centre WA 6983

PAGE 2
NO. 08-005511-1

and must be presented to an authorised officer of the Department upon request.

- 14 All holotypes and syntypes and a half share of paratypes of species or subspecies permitted to be permanently taken under this licence shall be donated to the Western Australian Museum. Duplicates (one pair in each case) of any species collected, which represents a significant extension of geographic range shall upon request be donated to the Western Australian Museum.
- 15 To prevent any unnecessary collecting in this State, all specimens and material taken and retained under the authority of this license shall, upon request, be loaned to the Western Australian Museum. Any unused portion or portions of any specimen collected under the authority of this license shall be offered to the Western Australian Museum for inclusion in its collection or made available to other scientific workers if so required.
- 16 Within one month of the expiration of this licence, the holder shall submit an electronic return into the department's Wildlife Licensing System, detailing the locality, site, geocode, date and number of each species of fauna captured, sighted or vouchered during the currency of the licence. A copy of any paper, report or thesis resulting from the research shall upon completion be lodged with the Director General.

Purpose

TFA 2324-0024 & BA27000875

Authorised Person

| Surname | Given name(s) |
|-----------|---------------|
| Walker | Christina |
| Crossley | Lachlan |
| Girando | Simon |
| Geidans | Lukas |
| Li | Yanlin |
| Berry | Lewis |
| Lohr | Michael |
| Hickling | Alex |
| Fintan | Angel |
| Constanza | Lutz |
| Hill | Chloe |

Date of Issue 10/07/2023
Valid From 10/07/2023
Date of Expiry 10/07/2023

Licensee: Mr Evan Nicholas Webb
Address PO Box 14
West Perth WA 6872
Australia

Issued by a Wildlife Licensing Officer of the Department of Parks and Wildlife under delegation from the Minister for Environment pursuant to section 133(1) of the Conservation and Land Management Act 1984.



Appendix D Fauna Desktop Assessment Results

Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd

SLR Project No.: 675.VX5890.00001

17 June 2024

Conservation Status:

State - Listed under Biodiversity Conservation Act 2016 or Department of Biodiversity, Conservation and Attractions Conservation,

Commonwealth - Listed under Environmental Protection and Biodiversity Conservation Act 1999.

CR - Critically Endangered, EN - Endangered, VU - Vulnerable, MI - Migratory, CD - Conservation Dependent fauna, OS - Other Specially Protected fauna, MA - Marine, P - Listed as Priority by DBCA.

Database:

NM - NatureMap, PMST - EPBC Protected Matters Search Tool, DBCA - DBCA Threatened and Priority Fauna database search, Field - Recorded during the current field survey.

Literature:

A - *Fauna Assessment Pinjarra - Williams Road (SLK 14 – 67)* (Harewood, 2017)

B - *Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Holyoake* (GHD, 2021a)

C - *Black Cockatoo Habitat Assessment – Pinjarra Racecourse* (Harewood, 2019)

D - *Chuditch Pre-mine Survey, Myara* (Way, 2013)

E - *Vertebrate Fauna Survey of Planned Mining Areas at Alcoa's Willowdale Larego Mining Region* (Environmental Management and Research Consultants, 2006)

F - *Terrestrial Fauna Survey and Black Cockatoo Habitat Assessment for Huntly Mine - Holyoake* (GHD, 2021a)

G - *Report for Rail Reserves in the Shire of Serpentine Jarrahdale - Spring Flora and Vegetation Survey and Fauna and Habitat Assessment* (GHD, 2012)

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | | | | | | | | | |
|-----------------|----------------------------------|----------------------|---------------------|---------------|----------|------|------|-------|------------|---|---|---|---|---|---|---|--|--|--|--|--|--|--|
| | | | State | Common wealth | NM | PMST | DBCA | | A | B | C | D | E | F | G | | | | | | | | |
| Amphibia | | | | | | | | | | | | | | | | | | | | | | | |
| Limnodynastidae | <i>Heleioporus albopunctatus</i> | Western Spotted Frog | | | | | | | | | | | | x | | | | | | | | | |
| | <i>Heleioporus eyrei</i> | Moaning Frog | | | | | | | | | x | | | x | x | | | | | | | | |
| | <i>Heleioporus inornatus</i> | Whooping Frog | | | | | | x | | | | | | x | | | | | | | | | |
| Myobatrachidae | <i>Crinia georgiana</i> | Quacking Frog | | | | | | x | x | x | | | | x | x | x | | | | | | | |
| | <i>Crinia glauerti</i> | Clicking Frog | | | | | | x | x | x | | | | x | | x | | | | | | | |
| | <i>Crinia insignifera</i> | Squelching Froglet | | | | | | | x | | | | | | | | | | | | | | |
| | <i>Crinia pseudinsignifera</i> | Bleating Froglet | | | | | | | x | x | | | | x | x | | | | | | | | |
| | <i>Geocrinia leai</i> | Ticking Frog | | | | | | | | x | | | | x | x | | | | | | | | |
| Pelodyadidae | <i>Litoria adelaidensis</i> | Slender Tree Frog | | | | | | | x | x | | | | x | x | | | | | | | | |
| | <i>Litoria moorei</i> | Motorbike Frog | | | | | | | | | x | | | | | | | | | | | | |

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | | | | | | | | | |
|--------------|--------------------------------|---|---------------------|---------------|----------|------|------|-------|------------|---|---|---|---|---|---|--|--|--|--|--|--|--|--|
| | | | State | Common wealth | NM | PMST | DBCA | | A | B | C | D | E | F | G | | | | | | | | |
| Birds | | | | | | | | | | | | | | | | | | | | | | | |
| Acanthizidae | <i>Acanthiza apicalis</i> | Inland Thornbill, Broad-tailed Thornbill | | | | | | | x | x | | | x | x | | | | | | | | | |
| | <i>Acanthiza chrysorrhoa</i> | Yellow-rumped Thornbill | | | | | | | x | x | | | x | x | | | | | | | | | |
| | <i>Acanthiza inornata</i> | Western Thornbill | | | | | | | x | | | | x | x | | | | | | | | | |
| | <i>Aphelocephala leucopsis</i> | Southern Whiteface | | VU | | x | | | | | | | | | | | | | | | | | |
| | <i>Gerygone fusca</i> | Western Gerygone | | | | | | x | x | | | | x | | | | | | | | | | |
| | <i>Sericornis maculatus</i> | Spotted Scrubwren | | | | | | x | x | x | | | x | x | | | | | | | | | |
| | <i>Smicronis brevirostris</i> | Weebill | | | | | | | x | x | | | x | | x | | | | | | | | |
| Accipitridae | <i>Accipiter cirrocephalus</i> | Collared Sparrowhawk | | | | | | | | x | | | x | | | | | | | | | | |
| | <i>Accipiter fasciatus</i> | Brown Goshawk | | MA | | | | | | | | | x | x | | | | | | | | | |
| | <i>Aquila audax</i> | Wedge-tailed Eagle | | | | | | | x | x | | | x | x | x | | | | | | | | |
| | <i>Elanus scriptus</i> | Letter-winged Kite | P4 | | | | | | | | | | | | x | | | | | | | | |
| | <i>Haliastur sphenurus</i> | Whistling Kite | | MA | | | | | | x | | | x | | x | | | | | | | | |
| | <i>Hieraaetus morphnoides</i> | Little Eagle | | | | | | | | | | | x | | | | | | | | | | |
| | <i>Lophoictinia isura</i> | Square-tailed Kite | | | | | | | | | | | x | | | | | | | | | | |
| | <i>Milvus migrans</i> | Black Kite | | | | | | | | | | | | | x | | | | | | | | |
| Aegothelidae | <i>Aegotheles cristatus</i> | Australian Owlet-nightjar | | | | | | | x | x | | | | x | | | | | | | | | |
| Alcedinidae | <i>Dacelo novaeguineae</i> | Laughing Kookaburra | | | | | | | x | x | | | x | x | x | | | | | | | | |
| | <i>Todiramphus sanctus</i> | Sacred Kingfisher | | MA | | | | | | x | | | x | x | x | | | | | | | | |
| Anatidae | <i>Anas superciliosa</i> | Pacific Black Duck | | | | | | | x | x | | | x | x | | | | | | | | | |
| | <i>Chenonetta jubata</i> | Australian Wood Duck, Wood Duck, Maned Duck | | | | | | | x | x | | | | x | | | | | | | | | |
| | <i>Cygnus atratus</i> | Black Swan | | | | | | | | x | | | | | | | | | | | | | |
| | <i>Oxyura australis</i> | Blue-billed Duck | P4 | | x | | x | | | | | | | | | | | | | | | | |
| | <i>Tadorna tadornoides</i> | Australian Shelduck, Mountain Duck | | | | | | | x | | | | x | | | | | | | | | | |

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | | |
|---------------|--------------------------------------|---------------------------|---------------------|---------------|----------|------|------|-------|------------|---|---|---|---|---|---|---|
| | | | State | Common wealth | NM | PMST | DBCA | | A | B | C | D | E | F | G | |
| | <i>Spilopelia senegalensis</i> | Laughing Turtle Dove | | | | | | | | | | | | | | x |
| Corvidae | <i>Corvus bennetti</i> | Little Crow | | | | | | | | | | | | | x | |
| | <i>Corvus coronoides</i> | Australian Raven | | | | | | | x | x | | | | x | | x |
| Cuculidae | <i>Cacomantis flabelliformis</i> | Fan-tailed Cuckoo | | MA | | | | | x | | | | | x | x | |
| | <i>Chalcites basalis</i> | Horsfield's Bronze Cuckoo | | MA | | | | | | x | | | | x | | |
| | <i>Chalcites lucidus</i> | Shining Bronze Cuckoo | | MA | | | | | | | | | | x | x | |
| Dicaeidae | <i>Dicaeum hirundinaceum</i> | Mistletoebird | | | | | | | | | | | | | x | |
| Estrilidae | <i>Stagonopleura oculata</i> | Red-eared Firetail | | | | | | | | | | | | x | x | |
| Falconidae | <i>Falco cenchroides</i> | Nankeen Kestrel | | MA | | | | | | | | | | | | x |
| | <i>Falco hypoleucos</i> | Grey Falcon | VU | VU | | x | | | | | | | | | | |
| | <i>Falco peregrinus</i> | Peregrine Falcon | OS | | x | | x | | | x | | | | | x | |
| Hirundinidae | <i>Hirundo neoxena</i> | Welcome Swallow | | MA | | | | | x | | | | | | | |
| | <i>Petrochelidon ariel</i> | Fairy Martin | | | | | | | | x | | | | | x | x |
| | <i>Petrochelidon nigricans</i> | Tree Martin | | MA | | | | | | x | | | | x | x | |
| Laridae | <i>Hydroprogne caspia</i> | Caspian Tern | MI | MI, MA | x | | x | | | | | | | | | |
| | <i>Sternula nereis nereis</i> | Fairy Tern | VU | VU | | x | | | | | | | | | | |
| Locustellidae | <i>Cincloramphus cruralis</i> | Brown Songlark | | | | | | | | x | | | | | x | |
| | <i>Cincloramphus mathewsi</i> | Rufous Songlark | | | | | | | | x | | | | | | |
| Maluridae | <i>Malurus elegans</i> | Red-winged Fairywren | | | | | | x | | x | | | | x | x | |
| | <i>Malurus leucopterus</i> | White-winged Fairywren | | | | | | | | x | | | | | | |
| | <i>Malurus pulcherrimus</i> | Blue-breasted Fairywren | | | | | | | | x | | | | | | |
| | <i>Malurus splendens</i> | Splendid Fairywren | | | | | | x | x | x | | | | x | x | x |
| | <i>Stipiturus malachurus</i> | Southern Emu-wren | | | | | | | | | | | | | x | |
| Megapodiidae | <i>Leipoa ocellata</i> | Malleefowl | VU | VU | x | x | x | | | | | | | | | |
| Meliphagidae | <i>Acanthorhynchus superciliosus</i> | Western Spinebill | | | | | | | x | x | | | | x | x | |
| | <i>Anthochaera carunculata</i> | Red Wattlebird | | | | | | | x | x | | | | x | x | x |

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | |
|-----------------|---|--|---------------------|---------------|----------|------|------|-------|------------|---|---|---|---|---|---|
| | | | State | Common wealth | NM | PMST | DBCA | | A | B | C | D | E | F | G |
| | <i>Anthochaera lunulata</i> | Western Little Wattlebird, Western Wattlebird | | | | | | | x | x | | | x | x | |
| | <i>Gavicalis virescens</i> | Singing Honeyeater | | | | | | | | | | | x | x | x |
| | <i>Lichmera indistincta</i> | Brown Honeyeater | | | | | | | x | x | | | x | x | x |
| | <i>Melithreptus brevirostris</i> | Brown-headed Honeyeater | | | | | | | | x | | | x | x | |
| | <i>Melithreptus chloropsis</i> | Western White-naped Honeyeater | | | | | | | x | x | | | x | x | |
| | <i>Phylidonyris niger</i> | White-cheeked Honeyeater | | | | | | | | x | | | x | | x |
| | <i>Phylidonyris novaehollandiae</i> | New Holland Honeyeater | | | | | | | x | x | | | x | x | x |
| | <i>Ptilotula ornata</i> | Yellow-plumed Honeyeater | | | | | | | | x | | | | | |
| Meropidae | <i>Merops ornatus</i> | Rainbow Bee-eater | | MA | | | | | | x | | | x | x | |
| Monarchidae | <i>Grallina cyanoleuca</i> | Magpie-lark | | MA | | | | | | x | | | | x | x |
| | <i>Myiagra inquieta</i> | Restless Flycatcher | | | | | | | | x | | | x | x | |
| Motacillidae | <i>Anthus australis australis</i> | Australian Pipit | | MA | | | | | | | | | x | | |
| | <i>Motacilla cinerea</i> | Grey Wagtail | MI | MI, MA | | x | | | | | | | | | |
| Neosittidae | <i>Daphoenositta chrysoptera</i> | Varied Sittella | | | | | | | | x | | | x | | |
| Pachycephalidae | <i>Colluricincla harmonica</i> | Grey Shrikethrush | | | | | | x | | x | | | x | | |
| | <i>Pachycephala fuliginosa occidentalis</i> | Western Golden Whistler, Western Whistler | | | | | | | x | x | | | x | x | |
| | <i>Pachycephala rufiventris</i> | Rufous Whistler | | | | | | | | | | | x | | x |
| Pandionidae | <i>Pandion haliaetus</i> | Osprey | MI | MI, MA | x | x | x | | | | | | | | |
| Pardalotidae | <i>Pardalotus punctatus</i> | Spotted Pardalote | | | | | | | x | | | | x | x | |
| | <i>Pardalotus striatus</i> | Striated Pardalote | | | | | | | x | | | | x | x | |

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | | |
|-------------------|--|---------------------------------------|---------------------|---------------|----------|------|------|-------|------------|---|---|---|---|---|---|---|
| | | | State | Common wealth | NM | PMST | DBCA | | A | B | C | D | E | F | G | |
| | <i>Calidris canutus</i> | Red Knot | EN | VU, MI, MA | | x | | | | | | | | | | |
| | <i>Calidris ferruginea</i> | Curlew Sandpiper | CR | CR, MI, MA | | x | | | | | | | | | | |
| | <i>Calidris melanotos</i> | Pectoral Sandpiper | MI | MI, MA | | x | | | | | | | | | | |
| | <i>Calidris ruficollis</i> | Red-necked Stint | MI | MI, MA | x | | x | | | | | | | | | |
| | <i>Numenius madagascariensis</i> | Far Eastern Curlew, Eastern Curlew | CR | CR, MI, MA | | x | | | | | | | | | | |
| | <i>Tringa glareola</i> | Wood Sandpiper | MI | MI, MA | x | | x | | | | | | | | | |
| | <i>Tringa nebularia</i> | Common Greenshank | MI | EN, MI, MA | x | x | x | | | | | | | | | |
| Strigidae | <i>Ninox boobook</i> | Boobook Owl | | MA | x | | x | x | | x | | | | | x | |
| Threskiornithidae | <i>Plegadis falcinellus</i> | Glossy Ibis | MI | MI, MA | x | | x | | | | | | | | | |
| | <i>Threskiornis spinicollis</i> | Straw-necked Ibis | | MA | | | | | | | | | | | x | x |
| Turnicidae | <i>Turnix variegatus</i> | Painted Button-quail | | | | | | | | x | | | x | x | | |
| Tytonidae | <i>Tyto javanica</i> | Eastern Barn Owl | | | | | | | | x | | | | | | |
| | <i>Tyto novaehollandiae</i> | Australian Masked Owl, Masked Owl | P3 | | x | | x | | | | | | | | x | |
| Zosteropidae | <i>Zosterops lateralis</i> | Grey-breasted White-eye, Silvereye | | MA | | | | | x | x | | | x | x | | |
| Mammalia | | | | | | | | | | | | | | | | |
| Burramyidae | <i>Cercartetus concinnus</i> | Western Pygmy-possum, Mundarda | | | | | | | | x | | | | | | |
| Canidae | <i>Canis familiaris</i> * | Dingo / Dog | | | | | | | | | | | | | | x |
| | <i>Vulpes vulpes</i> * | Red Fox | | | | | | | | | | | | | x | x |
| Dasyuridae | <i>Antechinus flavipes leucogaster</i> | Yellow-footed Antechinus, Mardo | | | | | | x | | x | | | | | x | |
| | <i>Dasyurus geoffroyi fortis</i> | Western Quoll, Chuditch | VU | VU | x | x | x | x | | x | | x | | x | | |
| | <i>Phascogale tapoatafa wambenger</i> | Wambenger Brush-tailed Phascogale | CD | | x | | x | | | | | | | | | |
| | <i>Sminthopsis gilberti</i> | Gilbert's Dunnart | | | | | | | | x | | | | | x | |
| Felidae | <i>Felis catus</i> * | Cat | | | | | | x | x | x | | | | | x | x |

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | |
|------------------|---|---------------------------------|---------------------|---------------|----------|------|-------|-------|------------|---|---|---|---|---|---|
| | | | State | Common wealth | NM | PMST | DBCAs | | A | B | C | D | E | F | G |
| Leporidae | <i>Oryctolagus cuniculus</i> * | Rabbit | | | | | | | | x | | | | x | x |
| Macropodidae | <i>Macropus fuliginosus melanops</i> | Western Grey Kangaroo | | | | | | x | x | x | | | | x | x |
| | <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | x | | x | x | x | x | | | | x | x |
| | <i>Setonix brachyurus</i> | Quokka | VU | VU | x | x | x | | | x | | | | x | |
| Molossidae | <i>Austronomus australis</i> | White-striped Free-tailed Bat | | | | | | | | x | | | | x | |
| | <i>Ozimops kitcheneri</i> | Western Free-tailed Bat | | | | | | | | x | | | | x | |
| Muridae | <i>Hydromys chrysogaster</i> | Water Rat, Rakali | P4 | | x | | x | x | | | | | | x | |
| | <i>Mus musculus</i> * | House Mouse | | | | | | | | | | | | x | |
| | <i>Rattus rattus</i> * | Black Rat | | | | | | x | | x | | | | x | |
| Myrmecobiidae | <i>Myrmecobius fasciatus fasciatus</i> | Numbat, Walpurti | EN | EN | x | x | x | | | | | | | | |
| Peramelidae | <i>Isoodon fusciventer</i> | Quenda | P4 | | x | | x | | | x | | | | x | x |
| Phalangeridae | <i>Trichosurus vulpecula hypoleucus</i> | Common Brushtail Possum, Koomal | | | | | | x | | x | | | | x | |
| Potoroidae | <i>Bettongia penicillata ogilbyi</i> | Brush-tailed Bettong, Woylie | CR | EN | x | x | x | | | | | | | | |
| Pseudocheiridae | <i>Pseudocheirus occidentalis</i> | Western Ringtail Possum | CR | CE | x | x | x | | | | | | | | |
| Suidae | <i>Sus scrofa</i> * | Pig | | | | | | x | x | x | | | | x | |
| Tachyglossidae | <i>Tachyglossus aculeatus acanthion</i> | Short-beaked Echidna | | | | | | x | x | x | | | | x | |
| Vespertilionidae | <i>Chalinolobus gouldii</i> | Gould's Wattled Bat | | | | | | | | x | | | | x | |
| | <i>Chalinolobus morio</i> | Chocolate Wattled Bat | | | | | | | | x | | | | x | |
| | <i>Falsistrellus mackenziei</i> | Western False Pipistrelle | P4 | | x | | x | | | x | | | | x | |
| | <i>Vespadelus regulus</i> | Southern Forest Bat | | | | | | | | x | | | | x | |

| Family | Scientific name | Common name | Conservation status | | Database | | | Field | Literature | | | | | | | | | | | | | | | |
|------------------|---------------------------------------|-------------------------------|---------------------|---------------|----------|------|------|-------|------------|---|---|---|---|---|---|---|--|--|--|--|--|--|--|--|
| | | | State | Common wealth | NM | PMST | DBCA | | A | B | C | D | E | F | G | | | | | | | | | |
| Reptiles | | | | | | | | | | | | | | | | | | | | | | | | |
| Agamidae | <i>Ctenophorus ornatus</i> | Ornate Crevice Dragon | | | | | | | | | | | | x | x | | | | | | | | | |
| | <i>Pogona minor</i> | Dwarf Bearded Dragon | | | | | | | | x | | | | x | x | x | | | | | | | | |
| Carphodactylidae | <i>Underwoodisaurus milii</i> | Southern Barking Gecko | | | | | | | | x | | | | x | | | | | | | | | | |
| Diplodactylidae | <i>Diplodactylus lateroides</i> | Speckled Stone Gecko | | | | | | | | | | | | | x | | | | | | | | | |
| | <i>Diplodactylus polyophthalmus</i> | Spotted Sandplain Gecko | | | | | | | | | | | | x | | | | | | | | | | |
| Elapidae | <i>Acanthophis antarcticus</i> | Southern Death Adder | P3 | | | | | | | | | | | | x | | | | | | | | | |
| | <i>Notechis scutatus</i> | Tiger Snake | | | | | | | | x | | | | x | x | | | | | | | | | |
| | <i>Pseudonaja affinis</i> | Dugite | | | | | | | | x | | | | x | | x | | | | | | | | |
| | <i>Suta nigriceps</i> | Mitchell's Short-tailed Snake | | | | | | | | x | | | | | x | | | | | | | | | |
| Gekkonidae | <i>Christinus marmoratus</i> | Marbled Gecko | | | | | | | | | | | | | x | | | | | | | | | |
| Pygopodidae | <i>Aprasia pulchella</i> | Pretty Worm-lizard | | | | | | | | | | | | x | x | | | | | | | | | |
| | <i>Delma fraseri</i> | Fraser's Delma | | | | | | | | x | | | | | | | | | | | | | | |
| Pythonidae | <i>Aspidites ramsayi</i> | Woma | P1 | | x | | x | | | | | | | | | | | | | | | | | |
| | <i>Morelia spilota</i> | Carpet Python | | | | | | | | x | | | | x | x | | | | | | | | | |
| Scincidae | <i>Acritoscincus trilineatus</i> | Western Three-lined Skink | | | | | | | | x | | | | x | x | | | | | | | | | |
| | <i>Cryptoblepharus buchananii</i> | Buchanan's Snake-eyed Skink | | | | | | | | | | | | | x | x | | | | | | | | |
| | <i>Cryptoblepharus plagiocephalus</i> | Péron's Snake-eyed Skink | | | | | | | | | | | | x | | | | | | | | | | |
| | <i>Ctenotus delli</i> | Dell's Ctenotus | P4 | | x | | x | | | | | | | x | | | | | | | | | | |
| | <i>Ctenotus labillardieri</i> | Common Southwest Ctenotus | | | | | | | | x | | | | x | x | | | | | | | | | |
| | <i>Ctenotus ora</i> | Coastal Plains Skink | P3 | | x | | x | | | | | | | | | | | | | | | | | |
| | <i>Egernia kingii</i> | King's Skink | | | | | | | | | | | | x | x | | | | | | | | | |



Appendix E Fauna Site Sheets

Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd

SLR Project No.: 675.VX5890.00001

17 June 2024

5870-CAM-01-LB

| | | | | | |
|--------------------------|-------------------|---|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 413199 | Northing | 6354177 |
| Landform and Soil | | | Rock | | |
| Landform | Mid slope | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sandy loam | | Surface stone size classes present | Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Black, Grey | | | | |
| Condition | | | Habitat Features | | |
| Quality | Very good | | Water Source | Absent | |
| Fire History | Burnt (1-5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | |
| Disturbance | None observed | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | | <i>Jarrah</i> | |
| Mid stratum | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | | |
| Ground stratum | Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | | <i>Sedges</i> | |



Fulcrum photo ID | 25fcd6ee-6219-42bf-b3be-55a6b20cf8de

5870-CAM-03-LB

| | | | | | |
|--------------------------|---------------------------------------|-------------------------------------|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 411051 | Northing | 6356534 |
| Landform and Soil | | | Rock | | |
| Landform | Mid slope | | Rock type/s | Laterite | |
| Aspect | North | | Surface stone cover | 5 - 25% | |
| Soil type | Sand | | Surface stone size classes present | Pebbles (<0.6 cm), Rocks (20 - 60 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Brown, Grey | | | | |
| Condition | | | Habitat Features | | |
| Quality | Disturbed | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | |
| Disturbance | Clearing | | | | |
| Introduced fauna | None observed | | Ground Cover | 51-75% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open forest (50-80%) | | <i>Jarrah and Stags</i> | |
| Mid stratum | Tall (>2 m) | Shrubland and/or heathland (50-80%) | | | |
| Ground stratum | Low (>0.5 m) | Hummock grassland (50-80%) | | <i>Sedges</i> | |



Fulcrum photo ID | fc74674b-03b7-4bd2-aa3a-62b70cf96373

5870-CAM-05-LB

| | | | |
|--|---|---------------------------------|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 412964 | Northing: 6354412 | |
| Landform and Soil | | Rock | |
| Landform: Undulating plain | Rock type/s: None | | |
| Aspect: Negligible | Surface stone cover: 0 - 5% | | |
| Soil type: Sand | Surface stone size classes present: | | |
| Soil colour: Black, Grey | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Present | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hummocks, Leaf litter, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah</i> | |
| Mid stratum: Tall (>2 m) | Closed shrubland and/or heathland (>80%) | | |
| Ground stratum: Mid (0.5-1 m) | Open rushland and/or sedgeland (20-50%) | <i>Sedges</i> | |



Fulcrum photo ID 17faecb2-102a-4ec6-aca-702f8a4624d4

5870-CAM-09-LB

| | | | |
|--|---|---------------------------------|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 413690 | Northing: 6350349 | |
| Landform and Soil | | Rock | |
| Landform: Undulating plain | Rock type/s: Laterite | | |
| Aspect: Negligible | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Rocks (6 - 20 cm) | | |
| Soil colour: Black | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Exfoliating rock, Hollows - logs, Hummocks, Leaf litter, Peeling bark, Termite mounds | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah</i> | |
| Mid stratum: Tall (>2 m) | Shrubland and/or heathland (50-80%) | | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Bankisa</i> | |



Fulcrum photo ID a75fcc57-c7ba-4ed5-b253-c5ce5fb5a3b3

5870-CAM-10-LB

| | | | |
|--|---|---------------------------------|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 413789 | Northing: 6348987 | |
| Landform and Soil | | Rock | |
| Landform: Upper slope | Rock type/s: Laterite | | |
| Aspect: East | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Big Rocks (60 cm - 2 m), Rocks (20 - 60 cm), Small Rocks (6 - 20 cm) | | |
| Soil colour: Black, Brown | | | |
| Condition | | Habitat Features | |
| Quality: Disturbed | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | | |
| Disturbance: Clearing | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah</i> | |
| Mid stratum: Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | |
| Ground stratum: Low (>0.5 m) | Sparse hummock grassland (0.25-20%) | <i>Pea shrubs</i> | |



Fulcrum photo ID b2e2a2fb-a23e-49c8-8755-c8e206abd71b

5870-CAM-12-LB

| | | | |
|--|---|---------------------------------|--|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 414687 | Northing: 6350647 | |
| Landform and Soil | | Rock | |
| Landform: Upper slope | Rock type/s: Laterite | | |
| Aspect: East | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Rocks (20 - 60 cm), Small Rocks (6 - 20 cm) | | |
| Soil colour: Black, Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Logs > 10 cm, Peeling bark, Rock crevices, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: Pig | Ground Cover: 11-25% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah</i> | |
| Mid stratum: Mid (1-2 m) | Shrubland and/or heathland (50-80%) | | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Grasses and sedges</i> | |



Fulcrum photo ID 2bf75440-826d-4cfd-bdc9-34bab94dc7be-401e4b8f-802f-4e93-

5870-CAM-13-LB

| | | | | | |
|--------------------------|---------------------------------------|--|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 413084 | Northing | 6352726 |
| Landform and Soil | | | Rock | | |
| Landform | Undulating plain | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Clay loam | | Surface stone size classes present | Pebbles (<0.6 cm) | |
| Soil colour | Orange | | | | |
| Condition | | | Habitat Features | | |
| Quality | High quality | | Water Source | Present | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | |
| Disturbance | None observed | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Woodland (20-50%) | | <i>Jarrah</i> | |
| Mid stratum | Tall (>2 m) | Closed shrubland and/or heathland (>80%) | | | |
| Ground stratum | Mid (0.5-1 m) | Open hummock grassland (20-50%) | | <i>Sedges</i> | |



Fulcrum photo ID 3ec77bc4-f51a-425c-8265-5a75050fb7de

5870-CAM-16-LB

| | | | | | |
|--------------------------|---------------------------------------|---|---|---|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-14 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 414500 | Northing | 6348638 |
| Landform and Soil | | | Rock | | |
| Landform | Mid slope | | Rock type/s | Laterite | |
| Aspect | North | | Surface stone cover | 5 - 25% | |
| Soil type | Sandy loam | | Surface stone size classes present | Pebbles (<0.6 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Black, Brown | | | | |
| Condition | | | Habitat Features | | |
| Quality | High quality | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | |
| Disturbance | None observed | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | | <i>Jarrah and Allocasuarina</i> | |
| Mid stratum | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | | |
| Ground stratum | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | | | |



Fulcrum photo ID 75423b83-ef94-4eb4-9337-cd370848f0ea

5870-CAM-17-LB

| | | | |
|--|---|--|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 412681 | Northing: 6352341 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | | Rock type/s: Laterite | |
| Aspect: South | | Surface stone cover: 25 - 50% | |
| Soil type: Sandy loam | | Surface stone size classes present: Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour: Black | | | |
| Condition | | Habitat Features | |
| Quality: Disturbed | | Water Source: Absent | |
| Fire History: Little or no fire evidence (>5 years) | | Microhabitats: Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | |
| Disturbance: Clearing | | | |
| Introduced fauna: None observed | | Ground Cover: 26-50% | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open woodland (0.25-20%) | <i>Jarrah</i> | |
| Mid stratum: Mid (1-2 m) | Open mallee shrubland (20-50%) | | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Small herbs and sedges</i> | |



Fulcrum photo ID: ced6cc7a-b41f-4505-bd5d-74128d1fc492

5870-CAM-18-LB

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 413797 | Northing: 6355266 | |
| Landform and Soil | | Rock | |
| Landform: Lower slope | | Rock type/s: Laterite | |
| Aspect: South | | Surface stone cover: 0 - 5% | |
| Soil type: Sandy loam | | Surface stone size classes present: Small Rocks (6 - 20 cm), Stones (2 - 6 cm) | |
| Soil colour: Black, Brown | | | |
| Condition | | Habitat Features | |
| Quality: Good | | Water Source: Absent | |
| Fire History: Burnt (1-5 years) | | Microhabitats: Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | |
| Disturbance: Clearing | | | |
| Introduced fauna: None observed | | Ground Cover: 11-25% | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open woodland (0.25-20%) | <i>Jarrah</i> | |
| Mid stratum: Mid (1-2 m) | Sparse shrubland and/or heathland (0.25-20%) | | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Grasses</i> | |



Fulcrum photo ID: 5afb0ba5-ce56-49ee-8750-62b00bf22720

5870-CAM-19-LB

| | | | | | |
|--------------------------|---------------------------------------|--|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-14 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 412917 | Northing | 6349007 |
| Landform and Soil | | | Rock | | |
| Landform | Upper slope | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sandy loam | | Surface stone size classes present | Pebbles (<0.6 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Brown, Grey | | | | |
| Condition | | | Habitat Features | | |
| Quality | Disturbed | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Logs > 10 cm, Peeling bark, Termite mounds | |
| Disturbance | Vehicle tracks | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Woodland (20-50%) | | <i>Jarrah</i> | |
| Mid stratum | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | | |
| Ground stratum | Low (>0.5 m) | Sparse hummock grassland (0.25-20%) | | <i>Grasses and sedges</i> | |



Fulcrum photo ID | 0e367c3b-216d-4f82-b937-7c26c7fa729e

5870-CAM-23-LB

| | | | | | |
|--------------------------|---------------------------------------|---|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 410877 | Northing | 6355590 |
| Landform and Soil | | | Rock | | |
| Landform | Undulating plain | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sandy loam | | Surface stone size classes present | Rocks (20 - 60 cm), Small Stones (0.6 - 2 cm) | |
| Soil colour | Black, Brown | | | | |
| Condition | | | Habitat Features | | |
| Quality | Disturbed | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | |
| Disturbance | Dieback | | | | |
| Introduced fauna | None observed | | Ground Cover | 51-75% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Woodland (20-50%) | | <i>Jarrah</i> | |
| Mid stratum | Tall (>2 m) | Shrubland and/or heathland (50-80%) | | | |
| Ground stratum | Mid (0.5-1 m) | Open rushland and/or sedgeland (20-50%) | | | |



Fulcrum photo ID | ad36fe10-d6cb-4c62-8e0a-bce88bd7e601

5870-CAM-24-LB

| | | | | | |
|--------------------------|---------------------------------------|---|---|---|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 410946 | Northing | 6355236 |
| Landform and Soil | | | Rock | | |
| Landform | Undulating plain | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sandy loam | | Surface stone size classes present | Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm) | |
| Soil colour | Black, Brown | | | | |
| Condition | | | Habitat Features | | |
| Quality | Disturbed | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hummocks, Leaf litter, Peeling bark, Woody debris | |
| Disturbance | Dieback | | | | |
| Introduced fauna | | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | | <i>Jarrah</i> | |
| Mid stratum | Tall (>2 m) | Shrubland and/or heathland (50-80%) | | | |
| Ground stratum | Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | | <i>Sedges</i> | |



Fulcrum photo ID c1050e10-ed05-42dd-b784-19e8fdb020af

5870-CAM-26-LB

| | | | | | |
|--------------------------|-------------------|---|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 412702 | Northing | 6355307 |
| Landform and Soil | | | Rock | | |
| Landform | Undulating plain | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sand | | Surface stone size classes present | Pebbles (<0.6 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Grey | | | | |
| Condition | | | Habitat Features | | |
| Quality | Very good | | Water Source | Absent | |
| Fire History | Burnt (1-5 years) | | Microhabitats | Hollows - logs, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | |
| Disturbance | Dieback | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Tall (>30 m) | Woodland (20-50%) | | <i>Jarrah</i> | |
| Mid stratum | Tall (>2 m) | Shrubland and/or heathland (50-80%) | | | |
| Ground stratum | Low (>0.5 m) | Open rushland and/or sedgeland (20-50%) | | <i>Sedges</i> | |



Fulcrum photo ID acb1d6d1-b690-4e55-b626-a5af26e4195e

5870-CAM-27-LB

| | | | | | |
|--------------------------|---------------------------------------|--|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-14 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 415029 | Northing | 6349231 |
| Landform and Soil | | | Rock | | |
| Landform | Undulating plain | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sandy loam | | Surface stone size classes present | Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Black, Brown | | | | |
| Condition | | | Habitat Features | | |
| Quality | High quality | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | |
| Disturbance | None observed | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Woodland (20-50%) | | <i>Jarrah</i> | |
| Mid stratum | Tall (>2 m) | Open shrubland and/or heathland (20-50%) | | | |
| Ground stratum | Low (>0.5 m) | Sparse hummock grassland (0.25-20%) | | <i>Xanthorrhoea</i> | |



Fulcrum photo ID c9935215-218e-4b14-a67f-d49c7360b426

5870-CAM-28-LB

| | | | | | |
|--------------------------|-------------------|---|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-13 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 412768 | Northing | 6354897 |
| Landform and Soil | | | Rock | | |
| Landform | Undulating plain | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 0 - 5% | |
| Soil type | Sand | | Surface stone size classes present | Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm) | |
| Soil colour | Grey | | | | |
| Condition | | | Habitat Features | | |
| Quality | High quality | | Water Source | Absent | |
| Fire History | Burnt (1-5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | |
| Disturbance | Dieback | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | | <i>Jarrah</i> | |
| Mid stratum | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | | |
| Ground stratum | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | | | |



Fulcrum photo ID 9922c67a-b31b-43a8-80b5-171f605037b6

5870-CAM-28-LB

| | | | |
|--|--|---|---------------|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 412768 | Northing: 6354897 | |
| Landform and Soil | | Rock | |
| Landform: Undulating plain | Rock type/s: Laterite | | |
| Aspect: Negligible | Surface stone cover: 0 - 5% | | |
| Soil type: Sand | Surface stone size classes present: Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm) | | |
| Soil colour: Grey | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | | |
| Disturbance: Dieback | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: | Mid (10-30 m) | Open woodland (0.25-20%) | <i>Jarrah</i> |
| Mid stratum: | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | |
| Ground stratum: | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | |



Fulcrum photo ID 9922c67a-b31b-43a8-80b5-171f605037b6

5870-CAM-30-LB

| | | | |
|--|---|--|---------------------------|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 414174 | Northing: 6352804 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Southeast | Surface stone cover: 5 - 25% | | |
| Soil type: Sandy loam | Surface stone size classes present: Pebbles (<0.6 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Black, Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: | Mid (10-30 m) | Open woodland (0.25-20%) | <i>Jarrah</i> |
| Mid stratum: | Mid (1-2 m) | Sparse shrubland and/or heathland (0.25-20%) | |
| Ground stratum: | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedges and grasses</i> |



Fulcrum photo ID fb7b0506-a3f8-46df-b841-382f3897ec83

5870-CAM-31-LB

| | | | |
|--|---|---------------------------------|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 411808 | Northing: 6355031 | |
| Landform and Soil | | Rock | |
| Landform: Undulating plain | Rock type/s: Laterite | | |
| Aspect: Negligible | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Pebbles (<0.6 cm), Rocks (20 - 60 cm), Small Stones (0.6 - 2 cm) | | |
| Soil colour: Grey | | | |
| Condition | | Habitat Features | |
| Quality: Disturbed | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | | |
| Disturbance: Dieback | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah</i> | |
| Mid stratum: Tall (>2 m) | Shrubland and/or heathland (50-80%) | | |
| Ground stratum: Mid (0.5-1 m) | Rushland and/or sedgeland (50-80%) | | |



Fulcrum photo ID /9c93bdb0-9d8c-4f70-b41e-4e17eed3a62d

5870-CAM-32-LB

| | | | |
|--|---|---------------------------------|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Basic Fauna | |
| Zone: 50 | Easting: 413233 | Northing: 6354593 | |
| Landform and Soil | | Rock | |
| Landform: Undulating plain | Rock type/s: Laterite | | |
| Aspect: Negligible | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Rocks (6 - 20 cm), Stones (2 - 6 cm) | | |
| Soil colour: Black | | | |
| Condition | | Habitat Features | |
| Quality: Disturbed | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Rock crevices, Woody debris | | |
| Disturbance: Dieback | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open woodland (0.25-20%) | <i>Jarrah</i> | |
| Mid stratum: Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | |
| Ground stratum: Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedges</i> | |



Fulcrum photo ID 9f907646-156e-4f71-bfeb-f9479c1f5d67

5870-CAM-34-LB

| | | | | | |
|--------------------------|---------------------------------------|--|---|--|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-14 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 412410 | Northing | 6352719 |
| Landform and Soil | | | Rock | | |
| Landform | Mid slope | | Rock type/s | Laterite | |
| Aspect | South | | Surface stone cover | 5 - 25% | |
| Soil type | Sandy loam | | Surface stone size classes present | Pebbles (<0.6 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour | Brown, Grey, Orange | | | | |
| Condition | | | Habitat Features | | |
| Quality | Disturbed | | Water Source | Absent | |
| Fire History | Little or no fire evidence (>5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Leaf litter, Peeling bark, Woody debris | |
| Disturbance | Clearing | | | | |
| Introduced fauna | None observed | | Ground Cover | 26-50% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | | <i>Jarrah and She-oak</i> | |
| Mid stratum | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | | | |
| Ground stratum | Mid (0.5-1 m) | Open rushland and/or sedgeland (20-50%) | | <i>Herbs</i> | |



Fulcrum photo ID 4511c0c9-29f2-4b05-8348-b92cb4a1981c

5870-CAM-35-LB

| | | | | | |
|--------------------------|-------------------|---|---|---|---------|
| Project: | 5890 | | | | |
| Date | 2023-10-15 | | Sample Type | Basic Fauna | |
| Zone | 50 | Easting | 414772 | Northing | 6352745 |
| Landform and Soil | | | Rock | | |
| Landform | Mid slope | | Rock type/s | Laterite | |
| Aspect | Negligible | | Surface stone cover | 25 - 50% | |
| Soil type | Sandy loam | | Surface stone size classes present | Rocks (20 - 60 cm), Small Rocks (6 - 20 cm) | |
| Soil colour | Black | | | | |
| Condition | | | Habitat Features | | |
| Quality | High quality | | Water Source | Absent | |
| Fire History | Burnt (1-5 years) | | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Logs > 10 cm, Peeling bark, Rock crevices, Woody debris | |
| Disturbance | None observed | | | | |
| Introduced fauna | None observed | | Ground Cover | 11-25% | |
| Vegetation | | | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | | <i>Jarrah</i> | |
| Mid stratum | Low (0.5-1 m) | Open shrubland and/or heathland (20-50%) | | <i>Jarrah</i> | |
| Ground stratum | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | | <i>Sedges and grasses</i> | |



Fulcrum photo ID 93f2c84d-146f-4179-94a7-40fbd1de8faa

5870-CAM-36-LB

| | | | |
|-------------------|--------------------------|-------------------------------------|--|
| Project: | | 5890 | |
| Date | | 2023-10-14 | |
| Sample Type | | Basic Fauna | |
| Zone | 50 | Easting | 412422 |
| Northing | | 6350668 | |
| Landform and Soil | | Rock | |
| Landform | Undulating plain | Rock type/s | None |
| Aspect | Negligible | Surface stone cover | 0 - 5% |
| Soil type | Sand | Surface stone size classes present | |
| Soil colour | Brown, Grey | | |
| Condition | | Habitat Features | |
| Quality | Disturbed | Water Source | Absent |
| Fire History | Burnt (1-5 years) | Microhabitats | Hollows - logs, Hollows - trees, Hummocks, Leaf litter, Peeling bark, Woody debris |
| Disturbance | Clearing, Vehicle tracks | Ground Cover | 26-50% |
| Introduced fauna | None observed | | |
| Vegetation | | | |
| Upper stratum | Mid (10-30 m) | Open woodland (0.25-20%) | <i>Jarrah</i> |
| Mid stratum | Mid (1-2 m) | Shrubland and/or heathland (50-80%) | |
| Ground stratum | Tall (1-2 m) | Sparse hummock grassland (0.25-20%) | <i>Herbs</i> |



Fulcrum photo ID [62a285cc-eb8a-4bcc-915d-c73916b8fd40](https://www.fulcrum.com.au/photo/62a285cc-eb8a-4bcc-915d-c73916b8fd40)

5890-CAM-02-EW

| | | | |
|-------------------|---------------------------------------|--|--|
| Project: | | 5890 | |
| Date | | 2023-10-15 | |
| Sample Type | | Terrestrial vertebrate fauna | |
| Zone | 50 | Easting | 414503 |
| Northing | | 6351029 | |
| Landform and Soil | | Rock | |
| Landform | Mid slope | Rock type/s | Laterite |
| Aspect | Southeast | Surface stone cover | 25 - 50% |
| Soil type | Sandy loam | Surface stone size classes present | Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) |
| Soil colour | Brown | | |
| Condition | | Habitat Features | |
| Quality | Good | Water Source | Absent |
| Fire History | Little or no fire evidence (>5 years) | Microhabitats | Leaf litter, Logs > 10 cm, Peeling bark, Rock crevices, Woody debris |
| Disturbance | Logging, Vehicle tracks | Ground Cover | 51-75% |
| Introduced fauna | None observed | | |
| Vegetation | | | |
| Upper stratum | Mid (10-30 m) | Woodland (20-50%) | <i>Marri, Allocasuarina, Jarrah, Banksia sessilis</i> |
| Mid stratum | Mid (1-2 m) | Sparse shrubland and/or heathland (0.25-20%) | <i>Bossiaea, Calytrix</i> |
| Ground stratum | Low (>0.5 m) | Sparse hummock grassland (0.25-20%) | <i>Sedge, herbs</i> |



Fulcrum photo ID [e4640a26-5456-4f3c-871d-2aba6ad8168c-5fd61c25-6187-](https://www.fulcrum.com.au/photo/e4640a26-5456-4f3c-871d-2aba6ad8168c-5fd61c25-6187-)

5890-CAM-04-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 413062 | Northing: 6352206 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: East | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Grey | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: Pig | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | | |
| Mid stratum: Tall (>2 m) | Closed shrubland and/or heathland (>80%) | <i>Bossiaea</i> , <i>Acacia pulchella</i> , <i>Bracken</i> | |
| Ground stratum: Absent | | | |



Fulcrum photo ID fa21d811-6c98-4246-a483-4b88583654fe

5890-CAM-06-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 410950 | Northing: 6354138 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Negligible | Surface stone cover: 50 - 75% | | |
| Soil type: Sandy loam | Surface stone size classes present: Rocks (20 - 60 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: 26-50% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open woodland (0.25-20%) | <i>Marr</i> , <i>Jarrah</i> , <i>Allocasuarina</i> | |
| Mid stratum: Low (0.5-1 m) | Open shrubland and/or heathland (20-50%) | <i>Calytrix</i> , <i>Bossiaea</i> | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | | |



Fulcrum photo ID e6648a78-370f-4442-9645-076a2df65829

5890-CAM-07-EW

| | | | |
|--|------------------------|---|---------------------------------------|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 411296 | Northing: 6350882 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | | Rock type/s: None | |
| Aspect: Northeast | | Surface stone cover: | |
| Soil type: Clay loam | | Surface stone size classes present: | |
| Soil colour: Black, Brown | | | |
| Condition | | Habitat Features | |
| Quality: Very good | | Water Source: Present | |
| Fire History: Little or no fire evidence (>5 years) | | Microhabitats: Leaf litter, Logs > 10 cm, Peeling bark, Termite mounds, Woody debris | |
| Disturbance: Pig rooting | | | |
| Introduced fauna: Pig | | Ground Cover: 76-100% | |
| Vegetation | | | |
| Upper stratum: | Mid (10-30 m) | Woodland (20-50%) | <i>Eucalyptus globulus, E. patens</i> |
| Mid stratum: | Tall (>2 m) | Closed shrubland and/or heathland (>80%) | <i>Myrtaceae, Bossiaea</i> |
| Ground stratum: | Tall (1-2 m) | Open rushland and/or sedgeland (20-50%) | <i>Sedge</i> |



5890-CAM-08-EW

| | | | |
|--|------------------------|---|----------------------------|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 411907 | Northing: 6352650 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | | Rock type/s: Laterite | |
| Aspect: West | | Surface stone cover: 0 - 5% | |
| Soil type: Loam | | Surface stone size classes present: Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | | Water Source: Present | |
| Fire History: Little or no fire evidence (>5 years) | | Microhabitats: Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | | Ground Cover: 76-100% | |
| Vegetation | | | |
| Upper stratum: | Mid (10-30 m) | Woodland (20-50%) | <i>Eucalyptus patens</i> |
| Mid stratum: | Tall (>2 m) | Closed shrubland and/or heathland (>80%) | <i>Bossiaea, Myrtaceae</i> |
| Ground stratum: | Tall (1-2 m) | Sparse rushland and/or sedgeland (0.25-20%) | |



5890-CAM-11-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 411883 | Northing: 6352707 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Southeast | Surface stone cover: 0 - 5% | | |
| Soil type: Loam | Surface stone size classes present: Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: Pig | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri</i> | |
| Mid stratum: Mid (1-2 m) | Shrubland and/or heathland (50-80%) | <i>Bossiaea, Macrozamia, Bracken, Tetratheca</i> | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, Tetratheca</i> | |



Fulcrum photo ID: b250b4f3-2ab4-4326-bc8b-f30d8c55ebc0

5890-CAM-14-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 415154 | Northing: 6353208 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: West | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: Potential pig sign | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri, Banksia grandis</i> | |
| Mid stratum: Tall (>2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea</i> | |
| Ground stratum: Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, Bracken, Tetratheca</i> | |



Fulcrum photo ID: 1ccce24f-c7ea-4019-88da-c958b3657bb0

5890-CAM-15-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 412292 | Northing: 6350989 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Southwest | Surface stone cover: 5 - 25% | | |
| Soil type: Loam | Surface stone size classes present: Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Black, Brown | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: None observed | Ground Cover: 51-75% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Marri, Jarrah</i> | |
| Mid stratum: Tall (>2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea, Bracken</i> | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedges, Tetratheca, Bracken</i> | |



5890-CAM-20-EW

| | | | |
|---|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 410834 | Northing: 6352823 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | Rock type/s: Laterite | | |
| Aspect: Northeast | Surface stone cover: 0 - 5% | | |
| Soil type: Loam | Surface stone size classes present: Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Present | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: Potential old pig sign | Ground Cover: 51-75% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri</i> | |
| Mid stratum: Tall (>2 m) | Shrubland and/or heathland (50-80%) | <i>Bossiaea, Bracken</i> | |
| Ground stratum: Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, Tetratheca</i> | |



5890-CAM-21-EW

| | | | |
|--|------------------------|--|---|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 411613 | Northing: 6353040 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | | Rock type/s: Laterite | |
| Aspect: Southeast | | Surface stone cover: 5 - 25% | |
| Soil type: Sandy loam | | Surface stone size classes present: Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: High quality | | Water Source: Present | |
| Fire History: Burnt (1-5 years) | | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | | Ground Cover: | |
| Vegetation | | | |
| Upper stratum: | Mid (10-30 m) | Woodland (20-50%) | <i>Marri, Jarrah</i> |
| Mid stratum: | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea, Xanthorrhoea, Macrozamia</i> |
| Ground stratum: | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | |



5890-CAM-22-EW

| | | | |
|--|------------------------|--|-----------------------------------|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 414325 | Northing: 6354625 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | | Rock type/s: Laterite | |
| Aspect: East | | Surface stone cover: 25 - 50% | |
| Soil type: Sandy loam | | Surface stone size classes present: Rocks (20 - 60 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | | Water Source: Absent | |
| Fire History: Burnt (1-5 years) | | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | | Ground Cover: 51-75% | |
| Vegetation | | | |
| Upper stratum: | Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah, Marri</i> |
| Mid stratum: | Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea, Bracken</i> |
| Ground stratum: | Low (>0.5 m) | Sparse forbland (0.25-20%) | <i>Stylidium, Boronia, Orchid</i> |



5890-CAM-29-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 412681 | Northing: 6351901 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Northeast | Surface stone cover: 0 - 5% | | |
| Soil type: Loam | Surface stone size classes present: Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: None observed | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri</i> | |
| Mid stratum: Mid (1-2 m) | Shrubland and/or heathland (50-80%) | <i>Bossiaea, Bracken</i> | |
| Ground stratum: Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | | |



5890-CAM-33-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 412669 | Northing: 6350011 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Southwest | Surface stone cover: 5 - 25% | | |
| Soil type: Loam | Surface stone size classes present: Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: None observed | Ground Cover: | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah, Marri</i> | |
| Mid stratum: Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea, Bracken</i> | |
| Ground stratum: Low (>0.5 m) | Sparse forbland (0.25-20%) | | |



5890-CAM-37-EW

| | | | |
|--|---|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: | Easting: | Northing: | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: South | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Burrows, Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri</i> | |
| Mid stratum: Tall (>2 m) | Closed shrubland and/or heathland (>80%) | <i>Bossiaea, Grevillea, Bracken</i> | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedges, Tetraheca</i> | |



5890-CAM-38-EW

| | | | |
|--|---|--|--|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 411260 | Northing: 6349796 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | Rock type/s: None | | |
| Aspect: East | Surface stone cover: | | |
| Soil type: Loam | Surface stone size classes present: | | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Eucalyptus globulus</i> | |
| Mid stratum: Tall (>2 m) | Closed shrubland and/or heathland (>80%) | <i>Myrtaceae, Bossiaea, Bracken, Lasioptalum</i> | |
| Ground stratum: Absent | | | |



5890-CAM-39-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 412821 | Northing: 6350751 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Northeast | Surface stone cover: 5 - 25% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah, Marri</i> | |
| Mid stratum: Tall (>2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea</i> | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, Tetratheca</i> | |



5890-CAM-40-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 412711 | Northing: 6353618 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: East | Surface stone cover: 0 - 5% | | |
| Soil type: Sandy loam | Surface stone size classes present: Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: None observed | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah, Marri, Banksia grandis</i> | |
| Mid stratum: Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea</i> | |
| Ground stratum: Mid (0.5-1 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, Xanthorrhoea</i> | |



5890-CAM-41-EW

| | | | |
|--|--|--|--|
| Project: 5890 | | | |
| Date: 2023-10-13 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 413985 | Northing: 6352460 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: Northwest | Surface stone cover: 25 - 50% | | |
| Soil type: Loam | Surface stone size classes present: Rocks (20 - 60 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: Pig | Ground Cover: 51-75% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri, Allocasuarina</i> | |
| Mid stratum: Tall (>2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea, Banksia grandis</i> | |
| Ground stratum: Mid (0.5-1 m) | Sparse tussock grassland (0.25-20%) | <i>Sedge, Tetratheca</i> | |



5890-CAM-42-EW

| | | | |
|--|---|--|--|
| Project: 5890 | | | |
| Date: 2023-10-14 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 413198 | Northing: 6351627 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | Rock type/s: None | | |
| Aspect: Northwest | Surface stone cover: | | |
| Soil type: Loam | Surface stone size classes present: | | |
| Soil colour: Brown | | | |
| Condition | | Habitat Features | |
| Quality: High quality | Water Source: Present | | |
| Fire History: Little or no fire evidence (>5 years) | Microhabitats: Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Open forest (50-80%) | <i>Eucalyptus patens</i> | |
| Mid stratum: Tall (>2 m) | Open shrubland and/or heathland (20-50%) | <i>Myrtaceae</i> | |
| Ground stratum: Tall (1-2 m) | Rushland and/or sedgeland (50-80%) | <i>Sedges, Xanthorrhoea, Lasiopetalum, Myrtaceae sp.</i> | |



5890-CAM-43-EW

| | | | |
|--|---|--|--|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 411849 | Northing: 6349686 | |
| Landform and Soil | | Rock | |
| Landform: Mid slope | Rock type/s: Laterite | | |
| Aspect: East | Surface stone cover: 25 - 50% | | |
| Soil type: Loam | Surface stone size classes present: Big Rocks (60 cm - 2 m), Rocks (20 - 60 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Absent | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: Logging | | | |
| Introduced fauna: None observed | Ground Cover: 51-75% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Jarrah, Marri, Allocasuarina</i> | |
| Mid stratum: Mid (1-2 m) | Open shrubland and/or heathland (20-50%) | <i>Bossiaea, Banksia grandis</i> | |
| Ground stratum: Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, prostrate Banksia</i> | |



5890-CAM-44-EW

| | | | |
|--|---|--|--|
| Project: 5890 | | | |
| Date: 2023-10-15 | | Sample Type: Terrestrial vertebrate fauna | |
| Zone: 50 | Easting: 414281 | Northing: 6351283 | |
| Landform and Soil | | Rock | |
| Landform: Drainage line | Rock type/s: None | | |
| Aspect: Southeast | Surface stone cover: | | |
| Soil type: Loam | Surface stone size classes present: | | |
| Soil colour: Brown, Orange | | | |
| Condition | | Habitat Features | |
| Quality: Very good | Water Source: Present | | |
| Fire History: Burnt (1-5 years) | Microhabitats: Hollows - logs, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris | | |
| Disturbance: None observed | | | |
| Introduced fauna: None observed | Ground Cover: 76-100% | | |
| Vegetation | | | |
| Upper stratum: Mid (10-30 m) | Woodland (20-50%) | <i>Eucalyptus globulus, Marri, Jarrah</i> | |
| Mid stratum: Tall (>2 m) | Shrubland and/or heathland (50-80%) | <i>Bossiaea, Myrtaceae</i> | |
| Ground stratum: Low (>0.5 m) | Isolated forbs (<0.25%) | | |



5890-CAM-45-EW

| | | | |
|--------------------------|---------------------------------------|---|---|
| Project: | 5890 | | |
| Date | 2023-10-14 | Sample Type | Terrestrial vertebrate fauna |
| Zone | 50 | Easting | 413041 |
| | | Northing | 6349511 |
| Landform and Soil | | Rock | |
| Landform | Mid slope | Rock type/s | Laterite |
| Aspect | Northeast | Surface stone cover | 0 - 5% |
| Soil type | Loam | Surface stone size classes present | Rocks (20 - 60 cm), Small Rocks (6 - 20 cm), Small Stones (0.6 - 2 cm), Stones (2 - 6 cm) |
| Soil colour | Brown | | |
| Condition | | Habitat Features | |
| Quality | Good | Water Source | Absent |
| Fire History | Little or no fire evidence (>5 years) | Microhabitats | Hollows - logs, Hollows - trees, Leaf litter, Logs > 10 cm, Peeling bark, Woody debris |
| Disturbance | Pig rooting | | |
| Introduced fauna | Pig | Ground Cover | 76-100% |
| Vegetation | | | |
| Upper stratum | Mid (10-30 m) | Open forest (50-80%) | <i>Jarrah, Marri</i> |
| Mid stratum | Mid (1-2 m) | Shrubland and/or heathland (50-80%) | <i>Bossiaea, Bracken, Grevillea</i> |
| Ground stratum | Low (>0.5 m) | Sparse rushland and/or sedgeland (0.25-20%) | <i>Sedge, Tetraetheca</i> |



Fulcrum photo ID d1930796-6ee4-4610-9652-e01e7800ec0



Appendix F Significant Fauna Recorded During the Survey

Targeted Chuditch, Quokka, and Woylie Survey

Willowdale Mine – Larego Region

Alcoa of Australia Ltd


SLR Project No.: 675.VX5890.00001

17 June 2024

| Scientific name | Common name | Conservation status | | Abundance | Date | Latitude | Longitude | Record type |
|-------------------------------------|----------------------------------|---------------------|---------------|-----------|------------|----------|-----------|---------------------|
| | | State | Common wealth | | | | | |
| <i>Zanda baudinii</i> | Baudin's Cockatoo | EN | EN | 1 | 13-10-2023 | -32.9519 | 116.0652 | Foraging evidence |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 1 | 13-10-2023 | -32.9519 | 116.0652 | Foraging evidence |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 1 | 14-10-2023 | -32.9706 | 116.0709 | Foraging evidence |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 1 | 14-10-2023 | -32.9606 | 116.0694 | Sighting |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 2 | 14-10-2023 | -32.9597 | 116.0817 | Feather |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 3 | 14-10-2023 | -32.9726 | 116.0779 | Sighting - overhead |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 6 | 15-10-2023 | -32.9744 | 116.0845 | Feather |
| <i>Calyptorhynchus banksii naso</i> | Forest Red-tailed Black Cockatoo | VU | VU | 6 | 15-10-2023 | -32.9609 | 116.0577 | Sighting |
| <i>Dasyurus geoffroii</i> | Chuditch | VU | VU | 1 | 18-10-2023 | -32.9758 | 116.0861 | Scat |
| <i>Hydromys chrysogaster</i> | Rakali | P4 | | 1 | 19-10-2023 | -32.9941 | 116.0524 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 14-10-2023 | -32.9480 | 116.0893 | Sighting |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 15-10-2023 | -32.9828 | 116.0641 | Sighting |

| Scientific name | Common name | Conservation status | | Abundance | Date | Latitude | Longitude | Record type |
|--------------------------|-----------------------|---------------------|---------------|-----------|------------|----------|-----------|-------------|
| | | State | Common wealth | | | | | |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 18-10-2023 | -32.9343 | 116.0467 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 18-10-2023 | -32.9877 | 116.0565 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 19-10-2023 | -32.9370 | 116.0662 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 20-10-2023 | -32.9593 | 116.0459 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 20-10-2023 | -32.9866 | 116.0502 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 23-10-2023 | -32.9734 | 116.0827 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 23-10-2023 | -32.9343 | 116.0467 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 24-10-2023 | -32.9768 | 116.0507 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 26-10-2023 | -32.9768 | 116.0507 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 26-10-2023 | -32.9939 | 116.0679 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 27-10-2023 | -32.9939 | 116.0679 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 27-10-2023 | -32.9792 | 116.0870 | Camera Trap |

| Scientific name | Common name | Conservation status | | Abundance | Date | Latitude | Longitude | Record type |
|--------------------------|-----------------------|---------------------|---------------|-----------|------------|----------|-----------|-------------|
| | | State | Common wealth | | | | | |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 29-10-2023 | -32.9973 | 116.0848 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 29-10-2023 | -32.9939 | 116.0679 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 30-10-2023 | -32.9451 | 116.0689 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 31-10-2023 | -32.9973 | 116.0848 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 01-11-2023 | -32.9768 | 116.0507 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 01-11-2023 | -32.9562 | 116.0922 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 2 | 03-11-2023 | -32.9734 | 116.0827 | Camera Trap |
| <i>Notamacropus irma</i> | Western Brush Wallaby | P4 | | 1 | 07-11-2023 | -32.9973 | 116.0848 | Camera Trap |



Making Sustainability Happen